

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 625 509 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
30.07.1997 Bulletin 1997/31

(21) Application number: **94303449.6**

(22) Date of filing: **13.05.1994**

(51) Int. Cl.⁶: **C07D 211/22, C07D 211/14,
C07D 211/26, C07D 211/28,
C07D 211/32, C07D 211/46,
A61K 31/445**

(54) N-alkyl substituted piperidine-derivatives with neurokinin receptor antagonist activity

N-alkylsubstituierte Piperidin-Derivate mit Neurokinin-Rezeptor antagonistischer Wirkung

Dérivés de pipéridine N-alkyl-substitués présentant une activité antagoniste des récepteurs de neurokinines

(84) Designated Contracting States:
CH DE FR GB IT LI
Designated Extension States:
SI

(30) Priority: **17.05.1993 GB 9310066**

(43) Date of publication of application:
23.11.1994 Bulletin 1994/47

(73) Proprietor: **ZENECA LIMITED**
London W1Y 6LN (GB)

(72) Inventors:
• **Jacobs, Robert Toms,**
ZENECA Inc.
Wilmington, Delaware 19897 (US)

• **Shenvi, Ashokkumar Bhikkappa,**
ZENECA Inc.
Wilmington, Delaware 19897 (US)

(74) Representative: **Smith, Stephen Collyer**
Intellectual Property Department
ZENECA Pharmaceuticals
Mereside
Alderley Park
Macclesfield Cheshire SK10 4TG (GB)

(56) References cited:
EP-A- 0 428 434

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 625 509 B1

Description

This invention concerns novel alkyl substituted heterocycles, and, more particularly, novel substituted piperidine derivatives which antagonize the pharmacological actions of one or more of the endogenous neuropeptide tachykinins known as neurokinins, particularly at the neurokinin 2 (NK2) receptor. The novel alkyl substituted heterocycles are useful whenever such antagonism is desired. Thus, such compounds may be of value in the treatment of those diseases in which an NK2 receptor is implicated, for example, in the treatment of asthma and related conditions. The invention also provides pharmaceutical compositions containing the novel alkyl substituted heterocycles for use in such treatment, methods for their use, and processes and intermediates for the manufacture of the novel alkyl substituted heterocycles.

The mammalian neurokinins comprise a class of peptide neurotransmitters which are found in the peripheral and central nervous systems. The three principal neurokinins are Substance P (SP), Neurokinin A (NKA) and Neurokinin B (NKB). There are also N-terminally extended forms of at least NKA. At least three receptor types are known for the three principal neurokinins. Based upon their relative selectivities favoring the neurokinin agonists SP, NKA and NKB, respectively, the receptors are classified as neurokinin 1 (NK1), neurokinin 2 (NK2) and neurokinin 3 (NK3) receptors, respectively. In the periphery, SP and NKA are localized in C-afferent sensory neurons, which neurons are characterized by non-myelinated nerve endings known as C-fibers, and are released by selective depolarization of these neurons, or selective stimulation of the C-fibers. C-Fibers are located in the airway epithelium, and the tachykinins are known to cause profound effects which clearly parallel many of the symptoms observed in asthmatics. The effects of release or introduction of tachykinins in mammalian airways include bronchoconstriction, increased microvascular permeability, vasodilation and activation of mast cells. Thus, the tachykinins are implicated in the pathophysiology and the airway hyperresponsiveness observed in asthmatics; and blockade of the action of released tachykinins may be useful in the treatment of asthma and related conditions. Peptidic NK2 antagonists have been reported. For example, a cyclic hexapeptide known as L-659,877 has been reported as a selective NK2 antagonist. Nonpeptidic NK2 antagonists also have been reported, for example in European Patent Application, Publication Number (EPA) 428434, EPA 474561, EPA 512901, EPA 512902 and EPA 515240, as well as in EPA 559538. We have discovered a series of nonpeptidic NK2 antagonists, and this is the basis for our invention.

According to the invention, there is provided a Compound of the invention which is a compound of formula I (formula set out hereinbelow following the Examples, together with other formulae denoted by Roman numerals) wherein

Q is phenyl which may bear one or two substituents independently selected from halo, trifluoromethyl, hydroxy, (1-3C)alkoxy, (1-3C)alkyl and methylenedioxy; or Q is thienyl, imidazolyl, benzo[b]thiophenyl or naphthyl any of which may bear a halo substituent; or Q is biphenyl; or Q is carbon-linked indolyl which may bear a benzyl substituent at the 1-position;

Q¹ is hydrogen or (1-3C)alkyl;

Q² is aryl or heteroaryl, which aryl or heteroaryl radical may bear one or more substituents independently selected from halo, hydroxy, (1-4C)alkoxy or (1-4C)alkyl;

R is (1-8C)alkyl or (3-8C)cycloalkyl which may bear one or more substituents selected from the group consisting of halo, (3-6C)cycloalkyl, cyano, nitro, hydroxy, lower alkoxy, lower acyloxy, aroyl, heteroaroyl, oxo, imino (which may bear a (1-6C)alkyl, (3-6C)cycloalkyl, (1-5C)acyl or aroyl substituent) hydroxyimino (which hydroxyimino may bear a lower alkyl or a phenyl substituent on the oxygen), an amino group of formula NR^AR^B, an amino group of formula NR^CR^D, an amidino group of formula C(=NR^G)NR^ER^F, and a carbamoyl group of formula CON(OR^K)R^L, but excluding any compound wherein a hydroxy and an oxo substituent together form a carboxy group, wherein

An amino group of formula NR^AR^B contains zero to about seven carbon atoms and each of R^A and R^B is independently hydrogen, (1-5C)alkyl or (3-6C)cycloalkyl, or R^A and R^B together with the nitrogen to which they are attached form a pyrrolidino, piperidino, morpholino, thiomorpholino (or its S-oxide) or piperazinyl group (which piperazinyl group may bear a methyl or ethyl group at the 4-position); and wherein R^C is hydrogen or lower alkyl and R^D is lower acyl, aroyl or heteroaroyl; or R^D is a group of formula C(=J)NR^ER^F in which J is oxygen, sulfur, NR^G or CHR^H; and wherein the amino group NR^ER^F contains zero to about seven carbon atoms and each of R^E and R^F is independently hydrogen, (1-5C)alkyl or (3-6C)cycloalkyl, or NR^ER^F together with the nitrogen to which they are attached form a pyrrolidino, piperidino, morpholino, thiomorpholino (or its S-oxide) or piperazinyl group (which piperazinyl group may bear a methyl or ethyl group at the 4-position) or R^E is hydrogen or lower alkyl and R^F together with R^G forms an ethylene or trimethylene group; R^G is hydrogen, lower alkyl or together with R^F forms an ethylene or trimethylene group; R^H is cyano, nitro or SO₂R^J and R^J is lower alkyl or phenyl; R^K and R^L are independently (1-3C)alkyl; and in which a cyclic group which is a substituent on R or formed by substitution on R may bear one or more (1-3C)alkyl groups on carbon as further substituents; and in which any aryl or heteroaryl group which is a part of the group R may bear one or more halo, lower alkyl, lower alkoxy, cyano, trifluoromethyl or nitro substituents; or a pharmaceutically acceptable salt thereof.

A particular subgroup of the invention is one in which Q is 3,4-dichlorophenyl, Q¹ is hydrogen and Q² is phenyl, or a pharmaceutically acceptable salt thereof.

It will be appreciated that a compound of formula I contains one or more asymmetrically substituted carbon atoms such that such compounds may be isolated in optically active, racemic and/or diastereomeric forms. Some compounds may exhibit polymorphism. It is to be understood that the present invention encompasses any racemic, optically-active, diastereomeric, polymorphic or stereoisomeric form, or mixtures thereof, which form possesses NK2 antagonist properties, it being well known in the art how to prepare optically-active forms (for example, by resolution of the racemic form or by synthesis from optically-active starting materials) and how to determine the NK2 antagonist properties by the standard tests described hereinafter. It may be preferred to use the compound of formula I in a form which is characterized as containing, for example, at least 95%, 98% or 99% enantiomeric excess of the form which corresponds to the (S)-configuration at the center indicated by * in formula I when Q is 3,4-dichlorophenyl, Q¹ is hydrogen and Q² is phenyl.

In this specification R, R^A, R^B, *et cetera* stand for generic radicals and have no other significance. It is to be understood that the generic term "(1-8C)alkyl" includes both straight and branched chain alkyl radicals but references to individual alkyl radicals such as "propyl" embrace only the straight chain ("normal") radical, branched chain isomers such as "isopropyl" being referred to specifically. A similar convention applies to other generic groups, for example, alkoxy, alkanoyl, *et cetera*. Halo is fluoro, chloro, bromo or iodo. Lower alkyl and lower alkoxy refer to radicals containing one to about four carbon atoms. Lower acyl and lower acyloxy refer to radicals containing one to about five carbon atoms. Aryl denotes a phenyl radical or an *ortho*-fused bicyclic carbocyclic radical having about nine to ten ring atoms in which at least one ring is aromatic. Heteroaryl encompasses a radical attached via a ring carbon of a monocyclic aromatic ring containing five or six ring atoms consisting of carbon and one to four heteroatoms selected from the group consisting of oxygen, sulfur and nitrogen, as well as a radical of an *ortho*-fused bicyclic heterocycle of about eight to ten ring atoms derived therefrom, particularly a benz-derivative or one derived by fusing a propenylene, trimethylene or tetramethylene diradical thereto, as well as a stable N-oxide thereof. Aryl and heteroaryl refer to arylcarbonyl and heteroarylcarbonyl radicals, respectively.

A pharmaceutically acceptable salt is one made with an acid which provides a physiologically acceptable anion.

Particular values are listed below for radicals, substituents and ranges for illustration only and they do not exclude other defined values or other values within defined ranges for the radicals and substituents. A particular value for Q is 3,4-dichlorophenyl, for Q¹ is hydrogen, and for Q² is phenyl. Particular values for (1-8C)alkyl include ethyl, propyl, butyl, pentyl, isopropyl, 3-pentyl and 4-heptyl. Particular values for a substituent on the (1-8C)alkyl group of R include hydroxy, acetoxy, oxo, hydroxyimino, methoxyimino, propylamino and acetamido. A particular value for two concurrent substituents on R is an oxo substituent, together on the same carbon with an amino group of formula NR^AR^B, thereby forming a carbamoyl group, for example an N-propylcarbamoyl group. Particular values for R include 1-hydroxyethyl, 1-hydroxypropyl, 1-hydroxybutyl, 1-hydroxy-1-methylethyl, 1-ethyl-1-hydroxy-propyl, 1-hydroxy-1-propylbutyl, 2-hydroxyethyl, acetyl, propionyl, 1-hydroxyiminoethyl, 1-hydroxyiminopropyl, 1-methoxyiminoethyl, 1-methoxyiminopropyl, 2-acetoxyethyl and 2-acetamidoethyl.

Specific compounds of formula I are described in the accompanying Examples. The compound of Example 16 is a preferred embodiment of the invention.

Pharmaceutically acceptable salts of a compound of formula I include those made with a strong inorganic or organic acid which affords a physiologically acceptable anion, such as, for example, hydrochloric, sulfuric, phosphoric, methanesulfonic, or *para*-toluenesulfonic acid.

A compound of formula I may be made by processes which include processes known in the chemical art for the production of structurally analogous heterocyclic compounds. Such processes and intermediates for the manufacture of a compound of formula I as defined above are provided as further features of the invention and are illustrated by the following procedures in which the meanings of generic radicals are as defined above unless otherwise indicated:

(a) Alkylating a corresponding piperidine of formula II with an aldehyde of formula III, by reductive alkylation, or with an alkylating agent of formula IV in which Y is a leaving group. The alkylation is preferably carried out by a conventional reductive alkylation, for example as described in Example 1, by the *in situ*, acid-catalyzed formation of an iminium salt, followed by reduction with sodium cyanoborohydride in alcoholic solvent. The reductive alkylation may be carried out in a suitable solvent, such as methanol, tetrahydrofuran or acidic water, using a suitable reducing agent, such as for example sodium cyanoborohydride, conveniently at a temperature in the range of -20 to 50 °C, preferably in the range of 0 to 25 °C.

(b) For a compound of formula I which is a ketone in which R bears an oxo substituent, condensation of a corresponding compound of formula V in which R¹ contains an N,N-disubstituted carbamoyl group, such as for example an N-methyl-N-methoxy carbamoyl group, with the required organomagnesium or organolithium reagent, such as for example an organomagnesium chloride or bromide (Grignard reagent). A Grignard addition may be carried out in a suitable solvent, such as for example diethyl ether or tetrahydrofuran, conveniently at a temperature in the range of -20 to 50 °C, preferably in the range of 0 to 25 °C. Suitable conditions for carrying out a Grignard addition can be found at Example 3.

(c) For a compound of formula I wherein R is a primary or secondary hydroxy, reduction of a corresponding compound of formula I wherein R is oxo, using a conventional reducing agent which does not reduce the benzamide moiety. Conveniently, the reduction may be carried out using sodium borohydride in a suitable solvent, such as for example methanol or ethanol, at a temperature in the range of -20 to 50 °C, preferably in the range of 0 to 25 °C. Suitable conditions for carrying out the reduction can be found at Example 4.

(d) For a compound of formula I wherein R is hydroxyimino (or an oxygen-substituted hydroxyimino), by condensation of a corresponding compound of formula I wherein R is oxo with hydroxylamine (or the corresponding oxygen-substituted hydroxylamine), conveniently as its hydrochloride salt. The condensation may be carried out in a suitable solvent, such as for example methanol or ethanol, and in the presence of a suitable base, such as for example pyridine. The reaction may conveniently be carried out at a temperature in the range of 0 to 100 °C, preferably at a temperature in the range of 25 to 75 °C. Suitable conditions for carrying out the condensation can be found at Example 5 or 6.

Whereafter, when it is desired to use a protecting group during all or a portion of an above described process; the protecting group then is removed when the final compound is to be formed.

Whereafter, for any of the above procedures, when a pharmaceutically acceptable salt of a compound of formula I is required, it may be obtained by reacting the compound of formula I with an acid affording a physiologically acceptable counterion or by any other conventional procedure.

If not commercially available, the necessary starting materials for the above procedures may be made by procedures which are selected from standard techniques of heterocyclic chemistry, techniques which are analogous to the synthesis of known, structurally similar compounds, and techniques which are analogous to the above described procedures or the procedures described in the Examples. The starting materials and the procedures for their preparation are additional aspects of the invention.

As will be clear to one skilled in the art, a variety of sequences is available for preparation of the starting materials, and the sequences leading to the starting materials and products of the invention may be altered if appropriate considerations regarding the synthetic methods and radicals present are followed.

An intermediate aldehyde of formula III may be prepared as outlined in Scheme I and described in Example 1, parts f.-i. for a compound in which Q is 3,4-dichlorophenyl, Q¹ is hydrogen and Q² is phenyl. Alkylation of the anion of 3,4-dichlorophenylacetonitrile with 1-bromo-2-(2-tetrahydropyranyloxy)ethane (conveniently prepared from 2-bromoethanol and dihydropyran using a strong acid exchange resin as catalyst) gives a nitrile of formula VI. Reduction of the nitrile gives a corresponding amine of formula VII which can be acylated using benzoic anhydride in the presence of a suitable base to give an amide of formula VIII. Alkylation of the amide with methyl iodide followed by hydrolysis of the acetal gives an alcohol of formula IX, which can be oxidized to give an intermediate of formula III. Alternatively, the alcohol of formula IX may be converted into an alkylating agent of formula IV using a conventional procedure.

An intermediate of formula III or of formula IV wherein the center marked with * has the (S) absolute configuration can be prepared from a corresponding compound of formula IX which may be obtained from a racemic compound of formula VII as outlined in Scheme II and described in Example 13, parts a.-f. for a compound in which Q is 3,4-dichlorophenyl, Q¹ is hydrogen and Q² is phenyl. Hydrolysis of an acetal of formula VII gives an amine of formula X. Salt formation with D-tartaric acid followed by crystallization, recrystallization and treatment with aqueous base gives the (S)-enantiomer of the compound of formula X. Treatment with ethyl chloroformate followed by reduction of the resulting carbamate gives the (S)-enantiomer of the amine of formula XI. Treatment of the amine with benzoyl chloride gives the (S)-enantiomer of the compound of formula IX, which can be oxidized to give the (S)-enantiomer of the compound of formula III or converted into the (S)-enantiomer of the compound of formula IV.

An intermediate amide of formula V may be prepared by alkylating a corresponding piperidine of formula IIa using a similar method to that described in (a) above or by any other conventional method.

The utility of a compound of the invention or a pharmaceutically acceptable salt thereof (hereinafter, collectively referred to as a "Compound") may be demonstrated by standard tests and clinical studies, including those disclosed in the EPA publications noted above, such as EPA 428434 or EPA 474561 (or US 5,236,921), and those described below.

Neurokinin A (NKA) Receptor-binding Assay (Test A)

The ability of a Compound of the invention to antagonize the binding of NKA at the NK2 receptor may be demonstrated using an assay using the human NK2 receptor expressed in Mouse Erythroleukemia (MEL) cells by using MEL cell membranes (MELM) which bear high-affinity and selective NK2 receptors and which is carried out as follows.

Brief Description of the Figures:

Figure 1 shows construction of the MEL cell expression vector construct pMEG3/hNK2R.

Figure 2 shows construction of the expression vector construct GSE1417/hNK2R.

Figure 3 shows expression of human NK2 receptor in MEL C88 cells.

MEL Cell Expression of Human NK2 Receptor (hNK2R):

Heterologous protein expression in Mouse Erythroleukemia (MEL) cells uses the human globin locus control region (LCR) (F. Grosveld et al., *Cell* (1987) 51, 975-985). The cDNAs are inserted between the human beta-globin promoter and the second intron of the human beta-globin gene, and this expression cassette is then placed downstream of the LCR and transfected into MEL cells (M. Needham et al., *Nucl. Acids Res.* (1992) 20, 997-1003). Human NK2 receptor cDNA (A. Graham et al., *Biochem. Biophys. Res. Commun.* (1991) 177, 8-16) was isolated from human lung RNA by polymerase chain reaction and DNA sequenced. Human NK2 receptor cDNA was subcloned into a shuttle vector (pMEG3) containing the beta-globin promoter and the 3' portion of the human beta-globin gene (Fig. 1). Human NK2 receptor cDNA was restricted with Eco 0109 (5' end) and Bam HI (3' end). An oligonucleotide linker-adaptor containing an internal Hind III site and a 3' end Eco 0109 site was ligated to the hNK2R cDNA fragment. The sequence of the top strand oligonucleotide = 5'd(GCGCAAGCTTATGGG) (SEQ ID NO:1) and the bottom strand oligonucleotide = 5'd(GTC-CCCATAAGCTTGCGC) (SEQ ID NO:2). These were annealed and ligated to the hNK2R fragment by standard methods. Following cleavage with Hind III, the resulting fragment was cloned into the Hind III and Bam HI sites in the polylinker of the shuttle vector pMEG3. The construct (pMEG3/hNK2R) was verified by restriction mapping and sequencing the 5' end and 3' end junctions of cDNA/vector. This was then transformed into *E. coli* DH5 alpha, and plasmid DNA was isolated by standard methods and verified by restriction mapping and DNA sequencing. A ClaI/Asp718 cassette carrying the beta-globin promoter, human NK2 receptor cDNA and the 3' beta-globin gene fragment was excised and subcloned downstream of the LCR in plasmid pGSE1417 (Fig. 2). The pMEG3/hNK2R construct was cleaved with ClaI and Asp718 and cloned directly into the ClaI and Asp718 sites (3' of LCR) in the expression vector GSE1417. The construct GSE1417/hNK2R (13.9kb) was verified by restriction mapping. *E. coli* DH5 alpha was transformed and recombinant plasmids verified by restriction mapping. MEL C88 cells (A. Deisseroth et al., *Cell* (1978) 15, 55-63) were electroporated (M. Antoniou, *Methods Molecular Biology* (1991) 7, 421-434) with PvuI linearized pGSE1417/human NK2 receptor DNA. Directly after transfection, cells were diluted in culture medium to 10^4 and 10^5 cell per mL and 1 mL aliquots transferred to each well of a 24-well plate. G418 was added to a concentration of 1 mg/mL 24 hours after the transfection to select for stable transfectants. Individual clones were picked or pooled to generate populations seven to ten days after the addition of selective medium. Fig. 3 shows the strategy used to isolate transfected MEL/human NK2 receptor cell line. For expression studies, cells were maintained in exponential growth for a period of four days, and then dimethyl sulfoxide (DMSO) was added to a final concentration of 2% (v/v) to induce differentiation and hence expression. Samples were taken 4 days post induction for mRNA and NKA binding analyses. The results indicated that clone #1 expresses hNK2R at the highest level (both hNK2R mRNA and specific NKA binding). This clone was scaled up and is now routinely fermented at 20 litre scale per month and supplied for use in Test A.

Membrane preparations (MELM) prepared from the MEL cells containing high-affinity NK2 receptors were prepared according to a published protocol (D. Aharony, et al., *Neuropeptides* (1992) 23, 121-130) with the following minor modifications: (1) Iodoacetamide (1 mM) was included in the homogenization buffer; (2) Homogenization was as published but for a shorter period of 10 seconds once and at a slower speed (setting 10); and (3) The equilibration step with KCl/EDTA was not performed. In a typical preparation, binding of ^3H -NKA (2.5 nM) to MELM was highly specific ($88 \pm 4\%$) and linearly dependent on the protein concentration, with significant binding detected as low as 26 μg protein/mL. Equilibrium-competition experiments demonstrated binding to high-affinity, high-density receptors with $K_D = 1187$ nM, $B_{\text{max}} = 2229$ fmol/mg protein.

The radio ligand ^3H -neurokinin A (^3H -NKA) as [4,5- ^3H -Leu 8]-NKA (typical specific activity, 117 Ci/mmol) is obtained by custom synthesis from Cambridge Research Biochemicals and is >95% pure. Repeated HPLC analysis demonstrated that the ligand is stable under proper storage conditions (silanized vials with 0.2% mercaptoethanol, under argon). Also, no degradation or metabolism is apparent in the receptor-binding assay.

The assay is carried out using an incubation buffer consisting of 50 mM Tris HCl (pH 7.4), 5 mM Mg^{++} , 100 μM thiorphan, 1 nM ^3H -NKA, 0.02% (w/v) BSA, 30 mM K^+ , and 300 μM dithiothreitol; and the concentration of membrane protein is held at approximately 0.05-0.025 mg per tube. Nonspecific binding is routinely defined with 1 μM NKA. Each tube receives the following: 150 μL incubation buffer, 20 μL ^3H -NKA, 20 μL Compound, NKA or buffer as appropriate, and 125 μL membrane suspension. The reaction is initiated by the addition of the membranes. The tubes are incubated for 60 min at 25 $^\circ\text{C}$ in a shaking water bath. The reaction is terminated by washing the tubes with 10 mL of ice-cold 50 mM Tris HCl using a Brandel cell harvesting system using Whatman GF/B filters which have been soaked at least 4 hours at room temperature in 0.01% (w/v) polyethylenimine to collect the membranes. The filters are deposited in scintillation vials and read in a Beckman LS 6000LL Scintillation Counter. The binding constant K_i is calculated by standard methods and is typically the mean of several such determinations. The K_i values may be converted to negative logarithms and expressed as $-\log$ molar K_i (i.e. $\text{p}K_i$).

In an initial use of this assay, the IC_{50} measured for the standard compound L-659,877 was found to be 30 nM versus ^3H -NKA binding to MELM. The selectivity of a Compound for binding at the NK2 receptor may be shown by deter-

mining its binding at other receptors using standard assays, for example, one using a tritiated derivative of SP in a tissue preparation selective for NK1 receptors or one using a tritiated derivative of NKB in a tissue preparation selective for NK3 receptors.

5 Guinea Pig Trachea Assay (Test B)

In the test described below either NKA or [β -ala⁸]-NKA(4-10) is used as an agonist. The chosen agonist is referred to as AG throughout the description. The ability of a Compound of the invention to antagonize the action of AG in a pulmonary tissue may be demonstrated using a functional assay in guinea pig trachea, which is carried out as follows.

10 Male guinea pigs are killed by a sharp blow to the back of the head. The trachea are removed, trimmed of excess tissue and divided into two segments. Each segment is suspended as a ring between stainless steel stirrups in water-jacketed (37.5 °C) tissue baths containing a physiological salt solution of the following composition (mM): NaCl, 119; KCl 4.6; CaCl₂, 1.8; MgCl₂, 0.5; NaH₂PO₄, 1; NaHCO₃, 25; glucose, 11; thiorphan, 0.001; and indomethacin, 0.005; gassed continuously with 95% O₂-5% CO₂. Initial tension placed on each tissue is 1 g, which is maintained throughout a 0.5 to 1.5 hour equilibration period before addition of other drugs. Contractile responses are measured on a Grass polygraph via Grass FT-03 force transducers.

Tissues are challenged repetitively with a single concentration of AG (10 nM) with intervening 30 min periods with washing to allow the tension to return to baseline levels. The magnitude of the contractions to AG reaches a constant level after two challenges, and each Compound is tested for inhibition of responses to AG by addition to the tissue bath 15 minute before the third or subsequent exposure to the agonist. The contractile response to AG in the presence of Compound is compared to that obtained with the second AG challenge (in the absence of Compound). Percent inhibition is determined when a Compound produces a statistically significant ($p < 0.05$) reduction of the contraction and is calculated using the second contractile response as 100%.

Potencies of selected Compounds are evaluated by calculating apparent dissociation constants (K_B) for each concentration tested using the standard equation:

$$K_B = [\text{agonist}] / (\text{dose ratio} - 1)$$

where dose ratio = $\text{antilog}[(\text{AG} - \log \text{molar EC}_{50} \text{ without Compound}) - (\text{AG} - \log \text{molar EC}_{50} \text{ with Compound})]$. The K_B values may be converted to the negative logarithms and expressed as $-\log \text{molar } K_B$ (i.e. pK_B). For this evaluation, complete concentration-response curves for AG are obtained in the absence and presence of Compound (30 min incubation period) using paired tracheal rings. The potency of AG is determined at 50% of its own maximum response level in each curve. The EC_{50} values are converted to the negative logarithms and expressed as $-\log \text{molar EC}_{50}$. Maximum contractile responses to AG are determined by expressing the maximum response to AG as a percentage of the contraction caused by carbachol (30 μM), added after the initial equilibration period. When a statistically significant ($p < 0.05$) reduction of the maximum response to AG is produced by a compound, the percent inhibition is calculated relative to the percentage of carbachol contraction in the untreated, paired tissue used as 100%.

40 Guinea Pig Labored Abdominal Breathing (Dyspnea) Assay (Test C)

Activity of a Compound of the invention as an antagonist of NKA at the NK2 receptor also may be demonstrated *in vivo* in laboratory animals, for example by adapting a routine guinea pig aerosol test described for evaluation of leukotriene antagonists by Snyder, et al. (Snyder, D. W., Liberati, N. J. and McCarthy, M. M., Conscious guinea-pig aerosol model for evaluation of peptide leukotriene antagonists. *J. Pharmacol. Meth.* (1988) 19, 219). Using the clear plastic chamber described previously by Snyder et al. to secure guinea pigs for a head-only aerosol exposure to bronchoconstrictor agonists, agonist is administered by aerosol to six conscious guinea pigs simultaneously during each maneuver. The tachykinin NK2-selective agonist, [β -ala⁸]-NKA(4-10), 3×10^{-5} M, is aerosolized from a Devilbiss Model 25 ultrasonic nebulizer into an air stream entering the chamber at a rate of 2 L/minute.

Guinea pigs (275 - 400 g) are fasted for approximately 16 hours prior to experimentation. Compounds to be evaluated for blockade of effects of [β -ala⁸]-NKA(4-10) or their vehicle (10% PEG400 in saline) are administered p.o. or i.v. at various times before aerosol agonist challenge. All animals are pretreated with atropine (10 mg/kg, i.p., 45 minutes pretreatment) indomethacin (10 mg/kg, i.p., 30 minutes pretreatment), propranolol (5 mg/kg, i.p., 30 minutes pretreatment), and thiorphan (1 mg/ml aerosol for 5 minutes, 15 minutes pretreatment).

Aerosol challenge with the agonist produces an initial increase in respiratory rate followed by a decrease with early signs of minor involvement of the abdominal muscles. The respiratory rate decreases further and the breathing becomes more labored with greater involvement of the abdominal muscles as exposure continues. The distinctly recognizable end point is the point where the breathing pattern of the guinea pig is consistently slow, deep, and deliberate, showing marked involvement of the abdominal muscles. Time, in seconds, from the onset of aerosol challenge to this end point is determined for each animal by using a stopwatch. The animals generally collapse after reaching the end

point and do not recover from the agonist-induced respiratory distress. Antagonists result in an increase in the time to reach the end point. Animals receive the aerosol administration of agonist for a maximum time of 780 seconds.

Differences between drug treated groups and corresponding vehicle treated control groups are compared using Student's t-test for unpaired observations.

5 Clinical studies to demonstrate the efficacy of a Compound of the invention may be carried out using standard methods. For example, the ability of a Compound to prevent or treat the symptoms of asthma or asthma-like conditions may be demonstrated using a challenge of inhaled cold air or allergen and evaluation by standard pulmonary measurements such as, for example, FEV₁ (forced expiratory volume in one second) and FVC (forced vital capacity), analyzed by standard methods of statistical analysis.

10 It will be appreciated that the implications of a Compound's activity in Test A or Test B is not limited to asthma, but rather, that the test provides evidence of general antagonism of NKA. In general, the Compounds of the invention which were tested demonstrated statistically significant activity in Test A with a K_i of 1 μ M or much less. For example, the compound described in Example 10 was typically found to have a K_i of 40 nM, and that of Example 16 a K_i of 1.5 nM. In Test B, a pK_B of 5 or greater was typically measured for a Compound of the invention. For example, a pK_B of 7.8 was measured for the compound described in Example 10, and a pK_B of 8.4 was measured for the compound described in Example 16. It should be noted that there may not always be a direct correlation between the activities of Compounds measured as K_i values in Test A and the values measured in other assays, such as the pK_B measured in Test B. No untoward side-effect following i.v. or p.o. dosing of a Compound of the invention in Test C has been noted. In Test C following oral administration at 5 μ mol/kg two hours prior to challenge with the agonist, the compound of Example 16 for example exhibited 100% protection, fully protecting all five of the animals tested.

20 As discussed above, a compound of formula I or a pharmaceutically acceptable salt thereof possesses NKA antagonist properties. Accordingly, it antagonizes at least one of the actions of NKA which are known to include bronchoconstriction, increased microvascular permeability, vasodilation and activation of mast cells. Accordingly, one feature of the invention is the use of a compound of formula I or a pharmaceutically acceptable salt thereof in the treatment of a disease in a human or other mammal in need thereof in which NKA is implicated and antagonism of its action is desired, such as, for example, the treatment of asthma or a related disorder. In addition, another feature of the invention is provided by the use of a compound of formula I or a salt thereof as a pharmacological standard for the development and standardization of new disease models or assays for use in developing new therapeutic agents for treating the diseases in which NKA is implicated or for assays for their diagnosis.

30 When used in the treatment of such a disease, a compound of the invention is generally administered as an appropriate pharmaceutical composition which comprises a compound of formula I or a pharmaceutically acceptable salt thereof as defined hereinbefore and a pharmaceutically acceptable diluent or carrier, the composition being adapted for the particular route of administration chosen. Such a composition is provided as a further feature of the invention. It may be obtained employing conventional procedures and excipients and binders, and it may be one of a variety of dosage forms. Such forms include, for example, tablets, capsules, solutions or suspensions for oral administration; suppositories for rectal administration; sterile solutions or suspensions for administration by intravenous or intramuscular infusion or injection; aerosols or nebulizer solutions or suspensions for administration by inhalation; or powders together with pharmaceutically acceptable solid diluents such as lactose for administration by insufflation.

40 For oral administration a tablet or capsule containing up to 250 mg (and typically 5 to 100 mg) of a compound of formula I may conveniently be used. For administration by inhalation, a compound of formula I will be administered to humans in a daily dose range of, for example, 5 to 100 mg, in a single dose or divided into two to four daily doses. Similarly, for intravenous or intramuscular injection or infusion a sterile solution or suspension containing up to 10% w/w (and typically 0.05 to 5% w/w) of a compound of formula I may conveniently be used.

45 The dose of a compound of formula I to be administered will necessarily be varied according to principles well known in the art taking account of the route of administration and the severity of the condition and the size and age of the patient under treatment. However, in general, the compound of formula I will be administered to a warm-blooded animal (such as man) so that a dose in the range of, for example, 0.01 to 25 mg/kg (and usually 0.1 to 5 mg/kg) is received. It will be understood that generally equivalent amounts of a pharmaceutically acceptable salt of a compound of formula I may be used.

50 The invention will now be illustrated by the following non-limiting examples in which, unless stated otherwise:

- (i) temperatures are given in degrees Celsius ($^{\circ}$ C); operations were carried out at room or ambient temperature, that is, at a temperature in the range of 18-25 $^{\circ}$ C;
- (ii) organic solutions were dried over anhydrous magnesium sulfate; evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000 pascals; 4.5-30 mm Hg) with a bath temperature of up to 60 $^{\circ}$ C;
- (iii) chromatography means flash chromatography on silica gel; reversed phase chromatography means chromatography over octadecylsilane (ODS) coated support having a particle diameter of 32-74 μ , known as "PREP-40-ODS" (Art 731740-100 from Bodman Chemicals, Aston, PA, USA); thin layer chromatography (TLC) was carried

out on silica gel plates;

(iv) in general, the course of reactions was followed by TLC and reaction times are given for illustration only;

(v) melting points are uncorrected and (dec) indicates decomposition; the melting points given are those obtained for the materials prepared as described; polymorphism may result in isolation of materials with different melting points in some preparations;

(vi) final products had satisfactory proton nuclear magnetic resonance (NMR) spectra;

(vii) yields are given for illustration only and are not necessarily those which may be obtained by diligent process development; preparations were repeated if more material was required;

(viii) when given, NMR data is in the form of delta values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as an internal standard, determined at 300 MHz using perdeuterio dimethyl sulfoxide (DMSO- d_6) as solvent; conventional abbreviations for signal shape are used; for AB spectra the directly observed shifts are reported; when given, coupling constants (J) are in Hz; Ar designates an aromatic proton when such an assignment is made;

(ix) chemical symbols have their usual meanings; SI units and symbols are used;

(x) reduced pressures are given as absolute pressures in pascals (Pa); elevated pressures are given as gauge pressures in bars;

(xi) solvent ratios are given in volume:volume (v/v) terms;

and (xii) mass spectra (MS) were run with an electron energy of 70 electron volts in the electron impact (EI) mode using a direct exposure probe; where indicated ionization was effected by chemical ionization (CI) or fast atom bombardment (FAB); values for m/z are given; generally, only ions which indicate the parent mass are reported.

Example 1. N-[2-(3,4-Dichlorophenyl)-4-[4-(1-hydroxyethyl)piperidino]butyl]-N-methylbenzamide hydrochloride.

A solution of 4-(1-hydroxyethyl)piperidine (450 mg) and N-[2-(3,4-dichlorophenyl)-3-formylpropyl]-N-methylbenzamide (1.18 g) in methanol (25 mL) was adjusted to pH 5.5 by addition of glacial acetic acid. This mixture was cooled to 0 °C and treated with sodium cyanoborohydride (304 mg). The resulting mixture was allowed to warm to room temperature and was stirred for 72 hours. The solvent was evaporated and the residue was dissolved in dichloromethane (100 mL) and washed successively with 1N aqueous sodium hydroxide, water (2x) and brine. The organic extract was dried, filtered and evaporated. Purification by chromatography (93:7:0.1 chloroform/methanol/aqueous ammonium hydroxide) afforded the pure title compound as a white foam (775 mg, 48%). Treatment of a solution of this foam (238 mg) in diethyl ether (5 mL) with hydrogen chloride (g) afforded the hydrochloride salt of the title compound (208 mg) as a white solid; mp 94-98 °C; NMR: 7.60-7.34 (m,5), 7.21-6.98 (m,3), 3.87-3.72 (m,2), 3.55 (br s,3), 3.28 (m,1), 3.15-2.77 (m,4), 3.07,2.76 (2s, NCH₃), 2.17-1.80 (br m,3), 1.54 (br m,4), 1.17 (d,3, J=6.2). MS (CI): m/z =465((M+1), ³⁷Cl), 463((M+1), ³⁵Cl). Analysis for C₂₅H₃₂Cl₂N₂O₂ • 1.0 HCl • 0.50 H₂O: Calculated: C, 59.00; H, 6.73; N, 5.50; Found: C, 58.98; H, 6.63; N, 5.44.

The starting material 4-(1-hydroxyethyl)piperidine was prepared as follows.

a. 1-(Benzyloxycarbonyl)-4-piperidine carboxylic acid. A solution of isonipecotic acid (5.0 g) in 10% aqueous sodium carbonate (100 mL) was cooled to 0 °C and treated with benzyl chloroformate (7.93 g). The resulting biphasic mixture was allowed to warm to room temperature over 2 hours, at which time it became homogeneous. The mixture was washed with ether (2x), was acidified to approximately pH 2 by addition of concentrated hydrochloric acid. The aqueous solution was extracted with ether (2x). The organic extracts were washed with brine, combined, dried, filtered and evaporated to leave a clear oil which slowly crystallized on standing (9.64 g); NMR (CDCl₃): 7.36 (m,5, C₆H₅), 5.13 (s,2, PhCH₂), 4.10 (br d,2, J=12.6 eq-H-C(2)), 2.96 (brdd,2, J=11.2, 12.6, ax-H-C(2)), 2.52 (m,1, H-C(4)), 1.93 (br d,2, J=10.8, eq-H-C(3)), 1.70 (m,2, ax-H-C(3)); MS (CI): m/z =264(M+1).

b. N-Methoxy-N-methyl-1-(Benzyloxycarbonyl)piperidine-4-carboxamide. A solution of 1-(benzyloxycarbonyl)-4-piperidinecarboxylic acid (2.0 g), N,N-dimethylhydroxylamine hydrochloride (890 mg), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.75 g), 4-dimethylaminopyridine (93 mg) and triethylamine (2.31 g) in dichloromethane (50 mL) was stirred at room temperature for 22 hours. The mixture was poured into 2N hydrochloric acid (50 mL) and extracted with ethyl acetate (3x). The organic extracts were washed sequentially with 2N hydrochloric acid, 10% aqueous sodium bicarbonate and brine, combined, dried, filtered and evaporated to afford the amide as a clear syrup (1.97 g); NMR (CDCl₃): d 7.35 (m,5, C₆H₅), 5.13 (s,2, PhCH₂), 4.25 (br,2), 3.71 (s,3, OCH₃), 3.19 (s,3, NCH₃), 2.86 (br,3), 1.71 (br,4).

c. 4-Acetyl-1-(benzyloxycarbonyl)piperidine. A solution of methyllithium (7 mL of a 1.4 M solution in diethyl ether) in tetrahydrofuran (15 mL) was cooled to -60 °C and treated dropwise with a solution of N-methoxy-N-methyl-1-(benzyloxycarbonyl)piperidine-4-carboxamide (2.50 g) in tetrahydrofuran (10 mL). The resulting solution was stirred at -60 °C for 2 hours, was warmed to 0 °C and quenched with 1N hydrochloric acid (10 mL). The mixture was diluted with 1N hydrochloric acid and extracted with ethyl acetate. The organic extract was washed sequentially with 1N hydrochloric acid and brine, dried, filtered and evaporated to afford a pale yellow syrup. Purification

by chromatography (dichloromethane) afforded the ketone as a white solid (1.12 g); NMR: 7.35 (m,5, C₆H₅), 5.06 (s,2, PhCH₂), 3.96 (m,2), 2.86 (m,2), 2.53 (m,1), 2.11 (s,3), 1.81 (m,2), 1.30 (m,2).

d. (RS)-1-Benzyloxycarbonyl-4-(1-hydroxyethyl)piperidine. A solution of 4-acetyl-1-(benzyloxycarbonyl)piperidine (980 mg) in methanol (25 mL) was treated with sodium borohydride (142 mg), added portionwise over 10 minutes. The resulting reaction mixture was stirred for 3 hours and evaporated. The residue was dissolved in dichloromethane (50 mL) and washed sequentially with water (2x) and brine. The organic extract was dried, filtered and evaporated to afford the alcohol as a clear syrup (959 mg); NMR: 7.35 (m,5, C₆H₅), 5.06 (s,2, PhCH₂), 4.03 (m,2), 3.54 (m,1), 2.70 (br m,2), 1.82 (m,1), 1.64 (m,1), 1.38 (m,1), 1.12 (m,2), 1.01 (d,3, J=6.3).

e. 4-(1-Hydroxyethyl)piperidine. A solution of 1-benzyloxycarbonyl-4-(1-hydroxyethyl)piperidine (959 mg) in ethanol (40 mL) was hydrogenated over 10% palladium on carbon (100 mg) at a hydrogen pressure of 3.45 bar for 16 hours. The resulting mixture was filtered through diatomaceous earth and evaporated to afford the piperidine as a clear oil (450 mg); NMR: 3.33 (m,1), 3.02 (br d,2, J=11.9), 2.46 (m,2), 1.73 (br d,1, J=12.7), 1.51 (br d,1, J=12.1), 1.26-1.03 (m,3), 1.00 (d,3, J=6.3).

The starting material N-[2-(3,4-dichlorophenyl)-3-formylpropyl]-N-methylbenzamide was prepared as follows.

f. 1-Bromo-2-(2-tetrahydropyranyloxy)ethane. To a mechanically stirred solution of dihydropyran (1000 mL) and Amberlyst 15 (10.0 g) in hexane (2000 mL) was added bromoethanol (985 g) dropwise over a period of 1.5 hours in a cold water bath to maintain an internal temperature of 35-40 °C. After being stirred overnight at room temperature, the reaction mixture was passed through a silica gel column (3 x 18 inches), and eluted with hexane (6 L). The eluate was evaporated to give an amber liquid which was distilled through a 2 inch vigreux collecting the material boiling between 75-95 °C at 3,300-4,700 Pa (25-35 mm Hg). This material was redistilled to give the ether (1195.5 g, 80-90 °C at 20 mm Hg) as an oil; NMR: 4.68 (m,1), 4.01 (m,1), 3.89 (m,1), 3.77 (m,1), 3.52 (m,3), 1.75-1.50 (m,6).

g. α-[2-(2-Tetrahydropyranyloxy)ethyl]-3,4-dichlorophenylacetonitrile. To a solution of sodium hydride (218.0 g of a 55% oil suspension) in tetrahydrofuran (4 L) at 10 °C in an ice/water bath was added 3,4-dichlorophenylacetonitrile (893.0 g) in tetrahydrofuran (2 L) over a period of 45 minutes, and the resulting solution was allowed to stir for 2 hours at room temperature. The mixture was cooled in an ice/water bath and 1-bromo-2-(2-tetrahydropyranyloxy)ethane (1076.0 g) was dropped in as a neat oil over a period of 25 minutes. The mixture was stirred overnight at room temperature and divided into four 2-liter portions. Each portion was diluted with saturated ammonium chloride (3 L) and extracted with ether (500 mL). The combined organic layers were washed (aqueous ammonium chloride), dried, and evaporated. The resulting material was chromatographed, with hexane/dichloromethane (gradient 100:0, 0:100) as eluent, to give the nitrile (932 g) as an oil; NMR: 7.47 (m,4), 7.20 (m,2), 4.57 (m,2), 4.08 (m,2), 3.85 (m,4), 3.54 (m,3), 3.37 (m,1), 2.15 (m,4), 1.77 (m,4), 1.56 (m,8).

h. 2-(3,4-Dichlorophenyl)-4-(2-tetrahydropyranyloxy)butylamine. To a solution of the above nitrile (128.3 g) in 95% ethanol (1.1 L) and concentrated ammonium hydroxide (550 mL) was added Raney Nickel (25.0 g). The mixture was hydrogenated under a hydrogen atmosphere at room temperature and 53 psi for 1.5 days. The mixture was filtered through diatomaceous earth to remove the catalyst, and the resulting filtrate was evaporated. The resulting material was chromatographed, with dichloromethane:methanol (gradient 100:0, 95:5) as eluent, to give the amine (91 g) as an oil; NMR: 7.40 (s,1), 7.38 (s,1), 7.32 (d,1, J=2.1), 7.28 (d,1, J=2.0), 7.07 (dd,1, J=2.1, 4.9), 7.04 (dd,1, J=2.1, 4.9), 4.50 (m,1), 4.43 (m,1), 3.70 (m,4), 3.45 (m,2), 3.27 (m,1), 3.17 (m,1), 2.97-2.75 (m,6), 2.00 (m,2), 1.82-1.66 (m,6), 1.53 (m,8), 1.18 (bs,4); MS (CI): m/z=318(M+1).

i. N-[2-(3,4-Dichlorophenyl)-4-(2-tetrahydropyranyloxy)butyl]-benzamide. To a solution of the above amine (2.5 g) in dichloromethane (35 mL) was added triethylamine (1.1 mL) and benzoic anhydride (1.85 g), and the resulting solution was allowed to stir for 45 minutes. The mixture was washed (0.2 N hydrochloric acid, 1 N sodium hydroxide, water), dried, and evaporated to give the amide (3.3 g) as an oil; NMR: 7.63 (m,4), 7.46 (m,2), 7.37 (m,8), 7.09 (m,2), 6.22 (m,2), 4.50 (m,1), 4.43 (m,1), 3.8 (m,5), 3.63 (m,1), 3.5 (m,4), 3.36 (m,1), 3.23 (m,1), 3.11 (m,2), 2.06 (m,2), 1.90-1.77 (m,4), 1.68 (m,2), 1.51 (m,8); MS (CI): m/z=338[(M+1)-tetrahydropyranyl].

j. N-[2-(3,4-Dichlorophenyl)-4-(2-tetrahydropyranyloxy)butyl]-N-methylbenzamide. To a solution of the above amide (3.3 g) in dimethyl sulfoxide (20 mL) was added powdered potassium hydroxide (1.6 g), followed by iodomethane (1.0 mL) after 15 minutes. After 1 hour, the mixture was diluted with water (330 mL) and extracted with dichloromethane (2 x 50 mL). The combined organic extracts were dried and evaporated to give the N-methyl-amide (3.1 g) as an oil; MS (CI): m/z=352[(M+1)-tetrahydropyranyl].

k. N-[2-(3,4-Dichlorophenyl)-4-hydroxybutyl]-N-methylbenzamide. To a solution of the above amide (10.5 g) in tetrahydrofuran (100 mL) was added 6 N hydrochloric acid (50 mL), and the resulting solution was allowed to stir overnight. The mixture was neutralized with 10 N sodium hydroxide, diluted with water (200 mL), and extracted with dichloromethane. The organic layer was dried and evaporated. The resulting yellow solid was suspended in ether and filtered to give the alcohol (8.4 g) as a white solid; MS (CI): m/z=352(M+1).

l. N-[2(3,4-Dichlorophenyl)-3-formylpropyl]-N-methylbenzamide. A solution of oxalyl chloride (878 mg) in dichloromethane (5 mL) was cooled to -78 °C and was treated dropwise with a solution of dimethyl sulfoxide (595 mg) in dichloromethane (2 mL). The resulting mixture was stirred at -78 °C for 15 minutes, and was treated dropwise with

EP 0 625 509 B1

a solution of *N*-[2-(3,4-dichlorophenyl)-4-hydroxybutyl]-*N*-methylbenzamide (1.22 g) in dichloromethane (10 mL)/dimethyl sulfoxide (2 mL). The mixture was stirred at -78 °C for 1 hour, was treated with triethylamine (1.75 g), warmed to room temperature and stirred for 1 hour. The mixture was poured into water and extracted with dichloromethane. The organic extract was washed sequentially with water (2x) and brine, dried, filtered through Florisil and evaporated to afford the aldehyde as a pale yellow oil (1.18 g); MS (CI): $m/z=352((M+1), ^{37}\text{Cl})$, $350((M+1), ^{35}\text{Cl})$.

Example 2. *N*-[2-(3,4-Dichlorophenyl)-4-[4-(1-hydroxybutyl)piperidino]butyl]-*N*-methylbenzamide hydrochloride.

Using a procedure similar to that described in Example 1, except using 4-(1-hydroxybutyl)piperidine, prepared by a similar sequence as described in Example 1(b-e), except using propylmagnesium chloride in place of methylolithium (step b), and the aldehyde prepared as described in Example 1.1., the hydrochloride of the title compound was obtained as a white powder, mp 92-95 °C; NMR: 7.60-7.36 (m,5), 7.20-6.97 (m,3), 3.82-3.76 (m,2), 3.58-3.47 (m,2), 3.36 (m,1), 3.25 (br s,1), 3.20-2.80 (m,4), 3.07, 2.75 (2s, NCH₃), 2.17-1.84 (br m,4), 1.67-1.30 (m,7), 0.94 (br s,3); MS (CI): $m/z=493((M+1), ^{37}\text{Cl})$, $491((M+1), ^{35}\text{Cl})$. Analysis for C₂₇H₃₆Cl₂N₂O₂ · 1.0 HCl · 1.0 H₂O: Calculated: C, 59.40; H, 7.20; N, 5.13; Found: C, 60.43; H, 7.05; N, 5.00.

Example 3. *N*-[2-(3,4-Dichlorophenyl)-4-(4-propionylpiperidino)butyl]-*N*-methylbenzamide hydrochloride.

A solution of ethylmagnesium chloride (0.47 mL of a 2M solution in tetrahydrofuran) in tetrahydrofuran (2 mL) was cooled to -78 °C and was treated dropwise with a solution of *N*-[2-(3,4-dichlorophenyl)-4-[4-(*N*-methoxy-*N*-methylcarbamoyl)piperidino]butyl]-*N*-methylbenzamide (400 mg) in tetrahydrofuran (5 mL). The resulting mixture was allowed to warm to room temperature and was stirred for 16 hours. The mixture was cooled to -78 °C and an additional quantity (0.47 mL) of ethylmagnesium chloride was added. The mixture was stirred for 2 hours at -78 °C, and was quenched with water and extracted with ethyl acetate. The organic extract was washed sequentially with water and brine, dried, filtered and evaporated to leave a pale yellow syrup. Purification by flash chromatography (95:5:0.1 dichloromethane/methanol/ammonium hydroxide) afforded the title compound as a white foam (217 mg). Treatment of this material (60 mg) with hydrogen chloride (g) in diethyl ether yielded the title compound as a white powder (47 mg), mp 87-90 °C; NMR: 7.59-7.32 (m,5), 7.20-6.97 (m,3), 3.84-3.74 (m,2), 3.70-3.48 (m,2), 3.13 (m,1), 3.05-2.76 (m,4), 3.06,2.75 (2s, NCH₃), 2.56 (q,2, J=7.2), 2.25-1.70 (br m,7), 1.01 (t,3, J=7.2); MS (CI): $m/z=477((M+1), ^{37}\text{Cl})$, $475((M+1), ^{35}\text{Cl})$. Analysis for C₂₆H₃₂Cl₂N₂O₂ · 1.0 HCl · 0.75 H₂O: Calculated: C, 59.43; H, 6.62; N, 5.33; Found: C, 59.50; H, 6.39; N, 5.33.

The starting material *N*-[2-(3,4-dichlorophenyl)-4-[4-(*N*-methoxy-*N*-methyl-carbamoyl)piperidino]butyl]-*N*-methylbenzamide was prepared as follows.

a. 4-(*N*-Methoxy-*N*-methylcarbamoyl)piperidine. A solution of *N*-methoxy-*N*-methyl-1-(benzyloxycarbonyl)piperidine-4-carboxamide (2.07 g), prepared as described in Example 1b, in ethanol (45 mL) was hydrogenated over 10% palladium on carbon (207 mg) at a hydrogen pressure of 3.45 bar for 4 hours. The mixture was filtered through diatomaceous earth and evaporated to afford the piperidine as a waxy, white solid (1.21 g); NMR: 3.67 (s,3, OCH₃), 3.08 (s,3, NCH₃), 2.96 (br m,2), 2.76 (br m,1), 2.50 (br m,2), 1.59-1.38 (m,4); MS (CI): $m/z=173(M+1)$.

b. *N*-[2-(3,4-Dichlorophenyl)-4-[4-(*N*-methoxy-*N*-methylcarbamoyl)piperidino]butyl]-*N*-methylbenzamide. A solution of 4-(*N*-methoxy-*N*-methylcarbamoyl)piperidine (500 mg) and *N*-[2-(3,4-dichlorophenyl)-3-formylpropyl]-*N*-methylbenzamide (885 mg), prepared as described in Example 1.1., in methanol (25 mL) was adjusted to pH 6 by addition of glacial acetic acid. The resulting mixture was cooled to 0 °C and treated with sodium cyanoborohydride (243 mg). The mixture was stirred for 0.5 hour at 0 °C, warmed to room temperature, stirred for 16 hours and evaporated. The residue was dissolved in dichloromethane, washed successively with 1N aqueous sodium hydroxide, water and brine, dried, filtered and evaporated to afford a colorless syrup (1.08 g). Purification by chromatography (93:7:1 dichloromethane/methanol/ammonium hydroxide) afforded *N*-[2-(3,4-dichlorophenyl)-4-[4-(*N*-methoxy-*N*-methylcarbamoyl)-piperidino]butyl]-*N*-methylbenzamide as a white foam (653 mg); NMR: 7.56-7.50 (m,4); 7.41-7.28 (m,4), 4.22 (s,1), 3.56 (s,3), 3.60-3.40 (m,4), 3.20 (s,3), 3.30-3.00 (m,2), 2.50 (m,1), 2.30 (m,1), 2.15 (m,1), 1.97 (m,2); MS: $m/z=451(M)$.

Example 4. *N*-[2-(3,4-Dichlorophenyl)-4-[4-(1-hydroxypropyl)-piperidino]butyl]-*N*-methylbenzamide hydrochloride.

A solution of *N*-[2-(3,4-dichlorophenyl)-4-[4-(propionyl)-piperidino]butyl]-*N*-methylbenzamide, prepared as described in Example 3, (148 mg) in methanol (10 mL) was cooled to 0 °C and treated with sodium borohydride (13 mg). The resulting mixture was stirred at 0 °C for 15 minutes, warmed to room temperature and stirred for 2 hours, and was evaporated. The residue was dissolved in dichloromethane (50 mL) and washed sequentially with water and brine. The organic extract was dried, filtered and evaporated to afford a yellow syrup (176 mg). Purification by chromatography (93:7:0.1 dichloromethane/methanol/ammonium hydroxide) afforded the title compound as a clear syrup (109 mg).

EP 0 625 509 B1

Treatment of this material with hydrogen chloride (g) in diethyl ether yielded the title compound as a white powder (92 mg), mp 95-97.5 °C; NMR: 7.60-7.33 (m,5), 7.20-6.97 (m,3), 3.84-3.76 (m,2), 3.57-3.48 (br m,2), 3.26 (m,2), 3.23-2.80 (m,4), 3.06,2.76 (2s, NCH₃), 2.30-1.78 (br m,4), 1.68-1.40 (m,5), 0.96 (t,3, J=7.2); MS (CI): m/z=479((H+1), ³⁷Cl), 477((M+1), ³⁵Cl). Analysis for C₂₆H₃₄Cl₂N₂O₂ · 1.0 HCl · 0.75 H₂O: Calculated: C, 59.21; H, 6.97; N, 5.31; Found: C, 59.05; H, 6.62; N, 5.62.

Example 5. N-[2-(3,4-Dichlorophenyl)-4-[4-((E)-1-oximinopropyl)-piperidino]butyl]-N-methylbenzamide.

A solution of N-[2-(3,4-dichlorophenyl)-4-[4-(propionyl)-piperidino]butyl]-N-methylbenzamide, prepared as described in Example 3, (200 mg), hydroxylamine hydrochloride (44 mg) and pyridine (166 mg) in ethanol (2 mL) was heated to reflux temperature and stirred for 3 hours. Upon cooling to room temperature, a white precipitate formed, which was isolated by filtration, washed with water and dried (150 °C/100 mbar) to afford the title compound (137 mg) as a white powder, mp 240-241.5 °C (dec); NMR (CD₃OD): 7.63-7.37 (m,5), 7.23-6.95 (m,3), 3.88-3.74 (m,2), 3.70-3.45 (m,2), 3.33-3.15 (m,3), 3.13-2.80 (m,4), 3.10,2.79 (2s, NCH₃), 2.22 (m,3), 2.03 (br m,4), 1.11 (t,3, J=7.1); MS (CI): m/z=492((M+1), ³⁷Cl), 490((M+1), ³⁵Cl). Analysis for C₂₆H₃₂Cl₂N₃O₂ · 2.25 H₂O: Calculated: C, 58.92; H, 6.94; N, 7.93; Found: C, 58.97; H, 6.93; N, 7.89.

Example 6. N-[2-(3,4-Dichlorophenyl)-4-[4-(1-((E)-methoximino)-propyl)piperidino]butyl]-N-methylbenzamide hydrochloride.

A solution of N-[2-(3,4-dichlorophenyl)-4-[4-(propionyl)-piperidino]butyl]-N-methylbenzamide, prepared as described in Example 3, (128 mg) and methoxylamine hydrochloride (34 mg) in pyridine (6 mL) was heated to reflux temperature and stirred for 6 hours. The mixture was evaporated, the residue suspended in water and extracted with dichloromethane. The organic extract was washed sequentially with water (2x) and brine, dried, filtered and evaporated to leave a tan foam. Purification by chromatography (95:5 dichloromethane/methanol) afforded a clear syrup, which was treated with hydrogen chloride (g) in diethyl ether to yield the title compound as a tan solid (46 mg), mp 83-85 °C; NMR: 7.61-7.36 (m,5), 7.21-6.95 (m,3), 3.83-3.71 (m,2), 3.77 (s, OCH₃), 3.70-3.45 (m,2), 3.29-2.71 (br m,5), 3.07,2.76 (2s, NCH₃), 2.48-1.72 (br m,9), 1.06 (t,3, J=7.4); MS (CI): m/z=506((M+1), ³⁷Cl), 504((M+1), ³⁵Cl). Analysis for C₂₇H₃₅Cl₂N₃O₂ · 1.0 HCl · 1.0 H₂O: Calculated: C, 58.02; H, 6.85; N, 7.52; Found: C, 58.17; H, 6.51; N, 7.42.

Example 7. N-[2-(3,4-Dichlorophenyl)-4-[4-(1-methyl-1-hydroxyethyl)-piperidino]butyl]-N-methylbenzamide hydrochloride.

Using a procedure similar to that described in Example 1, except using 4-(1-methyl-1-hydroxyethyl)piperidine, prepared as described below, and the aldehyde prepared as described in Example 1.1., the title compound was prepared and isolated as a white powder, mp 102-105 °C; NMR: 7.60-7.56 (m,2), 7.45-7.34 (m,3), 7.21 (m,2), 6.99 (m,1), 3.84-3.70 (m,2), 3.58 (m,2), 3.28 (m,1), 3.20-2.80 (m,4), 3.07,2.76 (2s, NCH₃), 2.20-1.90 (br m,4), 1.59 (br s,3), 1.18 (s,6); MS (CI): m/z=479((M+1), ³⁷Cl), 477((M+1), ³⁵Cl). Analysis for C₂₆H₃₄Cl₂N₂O₂ · 1.0 HCl · 0.50 H₂O: Calculated: C, 59.72; H, 6.94; N, 5.36; Found: C, 59.69; H, 6.87; N, 5.27.

The starting material 4-(1-hydroxy-1-methylethyl)piperidine was prepared as follows.

a. Ethyl 1-(benzyloxycarbonyl)-4-piperidinecarboxylate. A solution of ethyl 4-piperidinecarboxylate (14.71 g) and triethylamine (12.04 g) in chloroform (200 mL) was cooled to 0 °C and treated dropwise with benzyl chloroformate (17.91 g). The resulting mixture was stirred at 0 °C for 1 hour, allowed to warm to room temperature and was stirred for 12 hours. The mixture was washed with 1N hydrochloric acid (3x) and brine, dried, filtered and evaporated to yield a pale yellow oil (26.4 g). This material was purified by chromatography (95:5 dichloromethane/methanol) to afford the amide as a colorless syrup (22.15 g); NMR: 7.35 (m,5, C₆H₅), 5.07 (s,2, PhCH₂), 4.06 (q,2, J=7.0), 3.91 (br d,2, J=13.3), 2.95 (br,2), 2.53 (m,1), 1.82 (br d,2, J=13.3), 1.41 (m,2), 1.18 (t,3, J=7.0).

b. 1-(Benzyloxycarbonyl)-4-(1-hydroxy-1-methylethyl)piperidine. A solution of methyllithium (15.32 mL of a 1.4 M solution in diethyl ether) in tetrahydrofuran (15 mL) was cooled to -78 °C and treated dropwise with a solution of ethyl 1-(benzyloxycarbonyl)-4-piperidinecarboxylate (2.50 g) in tetrahydrofuran (10 mL). The resulting solution was stirred at -78 °C for 1 hour, warmed to 0 °C for 1 hour, quenched by addition of 1N hydrochloric acid and extracted with ethyl acetate. The organic extract was washed with 1N hydrochloric acid (2x) and brine, dried, filtered and evaporated to afford a pale yellow oil. Purification by chromatography (90:10 dichloromethane/methanol) gave the alcohol as a clear oil (940 mg); NMR: 7.35 (m,5, C₆H₅), 5.05 (s,2, PhCH₂), 4.04 (m,2), 2.69 (br,2), 1.68 (m,2), 1.35-1.03 (m,2), 1.01 (s,6); MS (CI): m/z=278(M+1).

c. 4-(1-Hydroxy-1-methylethyl)piperidine. A solution of 1-(benzyloxycarbonyl)-4-(1-hydroxy-1-methylethyl)piperidine (940 mg) in ethanol (30 mL) was hydrogenolysed over 10% palladium on carbon (100 mg) at a hydrogen pressure of 3.45 bar for 4 hours. The mixture was filtered and evaporated to afford the piperidine as a white foam (430

mg); NMR: 4.49 (s,1, OH), 2.97 (br d,2, J=11.7), 2.40 (m,2), 1.61 (br d,2, J=12.4), 1.27-1.03 (m,3), 1.00 (s,6).

Example 8. *N*-[2-(3,4-Dichlorophenyl)-4-[4-(1-ethyl-1-hydroxypropyl)piperidino]butyl]-*N*-methylbenzamide hydrochloride.

Using a procedure similar to that described in Example 1, except using 4-(1-ethyl-1-hydroxypropyl)piperidine, prepared by a route similar to that described in Example 7, except using ethylmagnesium chloride in place of methyllithium in step b, and the aldehyde prepared as described in Example 1.1., the title compound was isolated as a white solid, mp 102-104 °C; NMR: 7.59-7.32 (m,5), 7.20-6.97 (m,3), 3.83-3.74 (m,2), 3.56 (m,2), 3.28 (m,1), 3.25-2.80 (m,4), 3.20,2.75 (2s, NCH₃), 2.15 (br s,1), 1.90 (br s,3), 1.67-1.42 (m,7), 0.85 (t,6, J=7.3); MS (CI): m/z=507((M+1), ³⁷Cl), 505((M+1), ³⁵Cl). Analysis for C₂₈H₃₈Cl₂N₂O₂ • 1.0 HCl • 0.75 H₂O: Calculated: C, 60.54; H, 7.35; N, 5.04; Found: C, 60.33; H, 7.28; N, 4.98.

Example 9. *N*-[2-(3,4-Dichlorophenyl)-4-[4-(1-propyl-1-hydroxybutyl)piperidino]butyl]-*N*-methylbenzamide hydrochloride.

Using a procedure similar to that described in Example 1, except using 4-(1-propyl-1-hydroxybutyl)piperidine, prepared by a route similar to that described in Example 7, except using propylmagnesium chloride in place of methyllithium in step b, and the aldehyde prepared as described in Example 1.1., the title compound was isolated as a white solid; NMR: 7.60-7.32 (m,5), 7.20-6.97 (m,3), 3.85-3.70 (m,2), 3.60-3.48 (br m,2), 3.28 (m,1), 3.25-2.72 (m,4), 3.07,2.76 (2s, NCH₃), 2.17 (br s,1), 1.90 (br s,3), 1.66 (br s,3), 1.45-1.25 (m,8), 0.92 (t,6, J=7.1); MS (CI): m/z=535((M+1), ³⁷Cl), 533((M+1), ³⁵Cl). Analysis for C₃₀H₄₂Cl₂N₂O₂ • 1.0 HCl • 0.50 H₂O: Calculated: C, 62.23; H, 7.66; N, 4.84; Found: C, 61.98; H, 7.60; N, 4.72.

Example 10. *N*-[2-(3,4-Dichlorophenyl)-4-[4-(2-acetoxyethyl)piperidino]butyl]-*N*-methylbenzamide hydrochloride.

Using a procedure similar to that described in Example 1, except using 4-(2-acetoxyethyl)piperidine, prepared as described below, and the aldehyde prepared as described in Example 1.1., the title compound was isolated as a white solid; mp 62-71 °C; NMR: 7.8-6.9 (br,8), 4.02 (br,1), 1.99 (s,3); MS (CI): m/z=505(M+1). Analysis for C₂₇H₃₄Cl₂N₂O₃ • HCl • 1.5 H₂O: Calculated: C, 57.00; H, 6.73; N, 4.93; Found: C, 56.85; H, 6.60; N, 5.34.

The starting material 4-(2-acetoxyethyl)piperidine was prepared as follows.

a. 1-(Benzyloxycarbonyl)-4-(2-hydroxyethyl)piperidine. A solution of 4-(2-hydroxyethyl)piperidine (1.29 g) in dichloromethane (30 mL) was treated with triethylamine (1.02 g), cooled to 0 °C and treated dropwise with a solution of benzyl chloroformate (1.706 g) in dichloromethane (10 mL). The reaction mixture was allowed to warm to room temperature, stirred for 1 hour, diluted with dichloromethane (100 mL), and washed successively with 1N hydrochloric acid and brine. The aqueous layers were extracted with dichloromethane. The organic extracts were combined, dried, filtered and evaporated to afford the amide (2.27 g) as a clear amber oil; NMR: 7.35 (m,5), 5.06 (s,2), 4.71 (t,1, J=6), 4.00 (br,2), 3.42 (m,2), 2.7 (br,2), 1.6-1.0 (m,7); MS (CI): m/z=264(M+1).

b. 1-(Benzyloxycarbonyl)-4-(2-acetoxyethyl)piperidine. A solution of 1-benzyloxycarbonyl-4-(2-hydroxyethyl)piperidine (1.0 g) in dichloromethane (20 mL) was treated with triethylamine (0.38 g), cooled to 0 °C and treated dropwise with a solution acetyl chloride (0.3 g) in dichloromethane (5 mL). The reaction mixture was allowed to warm to room temperature and cooled back to 0 °C. Two additional portions of acetyl chloride (0.15 g each) were added and the reaction mixture was allowed to warm to room temperature. The mixture was stirred for 16 hours, diluted with dichloromethane (75 mL) and washed successively with 1N hydrochloric acid and brine. The aqueous washes were extracted with dichloromethane (50 mL). The organic extracts were combined, dried, filtered and evaporated to obtain the acetoxy compound (0.9 g) as a clear oil; NMR: 7.33 (br,5), 5.06 (s,2), 4.0 (br,4), 2.77 (br,2), 1.98 (s,3), 1.62 (d,1, J=13), 1.5-1.6 (m,4) 0.8-1.2 (m,2); MS (CI): m/z=306(M+1).

c. 4-(2-Acetoxyethyl)piperidine. A solution of 1-(benzyloxycarbonyl)-4-(2-acetoxyethyl)piperidine (0.9 g) in methanol (30 mL) was treated with a solution of anhydrous HCl in ether (1 mL) and was hydrogenated over 10% palladium on carbon at atmospheric pressure for 2 hours. The reaction mixture was filtered through diatomaceous earth and the filtrate was evaporated to afford the piperidine as a solid (0.60 g); NMR (d₆-dimethylsulfoxide with trifluoroacetic acid): 4.06 (t,3, J=6.5), 1.92 (s,3); MS (CI): m/z=172(M+1).

Example 11. *N*-[2-(3,4-Dichlorophenyl)-4-[4-(2-acetamidoethyl)piperidino]butyl]-*N*-methylbenzamide hydrochloride.

Using a procedure similar to that described in Example 1, except using 4-(2-acetamidoethyl)piperidine, prepared as described below, and the aldehyde prepared as described in Example 1.1., the title compound was isolated as a white solid; NMR: 7.8-6.9 (m,8), 3.30 (s,3), 3.28 (s,3); MS (CI): m/z=504(M+1). Analysis for C₂₇H₃₅Cl₂N₃O₂ • 1.0

HCl • 1.5 H₂O: Calculated: C, 56.65; H, 6.95; N, 7.34; Found: C, 56.62; H, 6.94; N, 7.29.

The starting material 4-(2-acetamidoethyl)piperidine was prepared as follows.

- a. 1-(Benzyloxycarbonyl)-4-(2-methanesulfonyloxyethyl)-piperidine. A solution of 1-benzyloxycarbonyl-4-(2-hydroxyethyl)piperidine, prepared as described in Example 10a, (0.79 g) in dichloromethane (10 mL) was treated with triethylamine (0.33 g), cooled to 0 °C, and treated with methanesulfonyl chloride (0.34 g). The reaction mixture was allowed to warm to room temperature, stirred for 2 hours, diluted with dichloromethane (50 mL), and washed successively with water and brine. The aqueous washes were extracted with dichloromethane. The combined organic extracts were dried, filtered and evaporated to afford the mesylate (0.97 g) as a clear syrup; NMR: 7.36 (br, s), 5.06 (s, 2), 4.24 (t, 2, J=6), 4.0 (m, 2), 3.16 (s, 3), 2.8 (br, 2), 1.5 (br, 3), 0.95-1.1 (m, 2); MS (FAB): 342(M+1).
- b. 1-(Benzyloxycarbonyl)-4-(2-azidoethyl)piperidine. A solution of the 1-(benzyloxycarbonyl)-4-(2-methanesulfonyloxyethyl)piperidine (0.97 g) in N,N-dimethylformamide (15 mL) was treated with sodium azide (0.37 g) and the resulting white suspension was stirred for 16 hours at the room temperature. The reaction mixture was diluted with ethyl acetate (100 mL) and washed with water (4x). The aqueous washes were extracted with ethyl acetate. The organic extracts were combined, dried, filtered and evaporated to afford the crude product (0.7 g). Purification by chromatography (9:1 hexane/ethyl acetate) afforded the azido compound as a clear oil (0.46 g); NMR: 7.35 (m, 5), 5.06 (s, 2), 4.01 (m, 2), 3.33-3.39 (m, 3), 2.78 (br, 2), 1.44-1.67 (m, 5), 1.00-1.05 (m, 2); MS (CI): m/z=289(M+1).
- c. 1-(Benzyloxycarbonyl)-4-(2-aminoethyl)piperidine. A solution of the 1-(benzyloxycarbonyl)-4-(2-azidoethyl)piperidine (0.46 g) in dry tetrahydrofuran (10 mL) was treated with triphenylphosphine (0.46 g). The resulting solution was stirred at 50 °C for 16 hours, was cooled to the room temperature, treated with aqueous ammonia (4.1 mL) and extracted with diethyl ether (3x). The organic extracts were combined and washed with 1N hydrochloric acid (3x). The combined aqueous extract was washed with ethyl acetate. The aqueous layer was made basic by addition of 50% aqueous sodium hydroxide, and was extracted with dichloromethane (3x). The organic extracts were combined, dried, filtered and evaporated to afford the amide as an oil (0.42 g); NMR: 7.34 (m, 5), 5.06 (s, 2), 4.0 (d, 2, J=13), 1.62 (d, 2, J=13), 1.55 (m, 1), 1.24-1.31 (m, 2), 0.95-1.00 (m, 2); MS (CI): m/z=263(M+1).
- d. 1-(Benzyloxycarbonyl)-4-(2-acetamidoethyl)piperidine. A solution of 1-(benzyloxycarbonyl)-4-(2-aminoethyl)piperidine (0.42 g) in dichloromethane (10 mL) was treated with triethylamine (0.16 g), cooled to 0 °C and treated with acetyl chloride (0.13 g), added in two portions. The reaction mixture was allowed to warm to room temperature and was stirred for 16 hours, was diluted with dichloromethane and washed successively with 1N hydrochloric acid and brine. The organic extract was dried, filtered and evaporated to afford a yellow solid (0.46 g). Purification by chromatography (95:5 dichloromethane:methanol) afforded 1-(benzyloxycarbonyl)-4-(2-acetamidoethyl)piperidine (0.27 g) as white solid; NMR: 7.8 (br, 1), 7.35 (m, 5), 5.06 (s, 2), 4.00 (d, 2, J=13), 3.04 (m, 2), 2.7 (br, 2), 1.78 (s, 3), 1.64 (d, 2, J=11), 1.35 (m, 1), 1.32 (m, 2), 0.99 (m, 2); MS (CI): m/z=305(M+1).
- e. 4-(2-Acetamidoethyl)piperidine. A solution of 1-(benzyloxycarbonyl)-4-(2-acetamidoethyl)piperidine (0.27 g) in methanol (10 mL) was hydrogenolysed over 10% palladium on carbon at atmospheric pressure for 16 hours. The mixture was filtered through diatomaceous earth and the filtrate was evaporated to afford the piperidine (0.13 g); NMR: 7.76 (br, 1), 3.02 (m, 2), 2.85 (m, 2), 2.38 (t, 2, J=11), 1.77 (s, 3), 1.54 (d, 2, J=12), 1.26 (m, 2), 0.96 (m, 2); MS (CI): m/z=171(M+1).

Example 12. N-[2-(3,4-Dichlorophenyl)-4-[4-(2-hydroxyethyl)piperidino]butyl]-N-methylbenzamide hydrochloride.

Using a procedure similar to that described in Example 1, except using 4-(2-hydroxyethyl)piperidine and the aldehyde prepared as described in Example 1.1., the title compound was isolated as a white solid, mp 45-52 °C; NMR (CD₃OD): 7.54-7.18 (m, 7), 6.99 (br, 1), 3.83-3.73 (m, 2), 3.63-3.45 (m, 4), 2.76 (s, 3), 2.17 (m, 1), 2.0-1.9 (m, 3), 1.71 (m, 1), 1.52-1.36 (m, 4); MS (CI): m/z=463(M+1). Analysis for C₂₅H₃₂Cl₂N₂O₂ • HCl • 1.5 H₂O: Calculated: C, 56.99; H, 6.87; N, 5.32; Found: C, 56.72; H, 6.31; N, 5.23.

Example 13. (S)-N-[2-(3,4-Dichlorophenyl)-4-(4-propionylpiperidino)butyl]-N-methylbenzamide hydrochloride.

- Using a procedure similar to that described in Example 3, except using (S)-N-[2-(3,4-dichlorophenyl)-4-[4-(N-methoxy-N-methylcarbamoyl)piperidino]butyl]-N-methylbenzamide, the title compound was isolated as a white solid, mp 85-90 °C; NMR: 7.60-7.34 (m, 5), 7.20-6.98 (m, 3), 3.85-3.72 (m, 2), 3.70-3.48 (m, 2), 3.26 (m, 1), 3.20-2.77 (m, 4), 3.07, 2.76 (2s, NCH₃), 2.57 (q, 2, J=7.1), 2.25-1.70 (br m, 7), 1.02 (t, 3, J=7.2); MS (CI): m/z=477((M+1), ³⁷Cl), 475((M+1) ³⁵Cl). Analysis for C₂₆H₃₂Cl₂N₂O₂ • 1.0 HCl • 0.75 H₂O: Calculated: C, 59.43; H, 6.62; N, 5.33; Found: C, 59.58; H, 6.33; N, 5.17.

The starting material (S)-N-[2-(3,4-dichlorophenyl)-4-[4-(N-methoxy-N-methylcarbamoyl)piperidino]butyl]-N-methylbenzamide was prepared as follows.

- a. 2-(3,4-Dichlorophenyl)-4-hydroxybutylamine. A solution of 2-(3,4-dichlorophenyl)-4-(2-tetrahydropyrany-

loxy)butylamine (550 g), prepared as described in Example 1.h., in methanol (3.3 L) was treated with 6N hydrochloric acid (352 mL). The reaction mixture was stirred for 3 hours and evaporated. The residue was dissolved in water (3 L) and extracted with diethyl ether. The aqueous solution was made basic by addition of sodium hydroxide, and extracted with ethyl acetate (4x). The organic extracts were combined, washed with brine, dried and evaporated to afford the alcohol as a waxy yellow solid (367 g); NMR (CDCl₃): 7.38 (d, 1, J=8.3), 7.28 (d, 1, J=2.1); 7.03 (dd, 1, J=2.1, 8.3), 3.66 (br, 1), 3.52 (br, 1), 3.1-2.3 (br, 5), 1.91 (br, 2).

b. (S)-2-(3,4-Dichlorophenyl)-4-hydroxybutylamine. A solution of D-tartaric acid (77.85 g) in methanol (2 L) was heated to reflux temperature and was treated with a solution of 2-(3,4-dichlorophenyl)-4-hydroxybutylamine (120.0 g) in methanol (0.8 L). The resulting mixture was stirred at reflux for 0.5 hour and was allowed to slowly cool to room temperature. The crystals which formed on standing (60.8 g) were isolated by filtration and dried. Concentration of the mother liquor to ca. 1 L afforded an additional quantity of salt (12.3 g), which was isolated by filtration and dried. The combined tartrate salt (73.1 g) was recrystallized from methanol (3.7 L), to afford the optically enriched salt (49.85 g). The salt was suspended in water (200 mL), treated with 1N aqueous sodium hydroxide (500 mL) and extracted with dichloromethane (5x). The dichloromethane extracts were combined, washed with brine, dried, filtered and evaporated to afford the optically enriched alcohol (30.6 g) as a waxy white solid; NMR (CD₃OD): 7.46 (d, 1, J=8.3), 7.42 (d, 1, J=2.1), 7.18 (dd, 1, J=2.1, 8.3), 4.84 (brs, 2), 3.50-3.30 (m, 2), 2.87-2.78 (m, 3), 1.93 (m, 1), 1.73 (m, 1); MS (CI): m/z=238((M+1), ³⁷Cl ³⁷Cl), 236((M+1), ³⁷Cl ³⁵Cl), 234((M+1), ³⁵Cl ³⁵Cl).

c. Ethyl (S)-N-[2-(3,4-dichlorophenyl)-4-hydroxybutyl]-carbamate. A solution of (S)-2-(3,4-dichlorophenyl)-4-hydroxybutylamine (30.4 g) and triethylamine (15.2 g) in dichloromethane (400 mL) was cooled to -25 °C and treated dropwise with a solution of ethyl chloroformate (15.2 g) in dichloromethane (400 mL). The mixture was allowed to warm to room temperature and stirred for 16 hours. The reaction was washed with 1N hydrochloric acid, 10% aqueous sodium bicarbonate and brine, dried, filtered and evaporated to afford an amber syrup. Purification by chromatography (1:1 dichloromethane:ethyl acetate) gave the amide (28.95 g) as a light yellow oil; NMR (CDCl₃): 7.42 (d, 1, J=8.3), 7.29 (d, 1, J=2.1), 7.05 (dd, 1, J=2.1, 8.3), 4.74 (br, 1), 4.10 (m, 2); 3.65-3.46 (m, 3), 3.25 (m, 1), 2.98 (m, 1), 1.89 (m, 1), 1.76 (m, 1), 1.23 (m, 3); MS (CI): m/z=310((M+1), ³⁷Cl ³⁷Cl), 308((M+1), ³⁷Cl ³⁵Cl), 306((M+1), ³⁵Cl ³⁵Cl).

d. (S)-N-[2-(3,4-Dichlorophenyl)-4-hydroxybutyl]-N-methylamine. A solution of ethyl (S)-N-[2-(3,4-dichlorophenyl)-4-hydroxybutyl]carbamate (28.9 g) in tetrahydrofuran (200 mL) was added dropwise to a suspension of lithium aluminum hydride (7.3 g) in tetrahydrofuran (100 mL). The reaction mixture was heated to reflux for 2 hours, cooled to 0 °C, and quenched by dropwise addition of saturated aqueous sodium sulfate solution (25 mL). The suspension was filtered through diatomaceous earth, the filter cake washed with tetrahydrofuran (6x) and the filtrate evaporated to yield (S)-N-[2-(3,4-dichlorophenyl)-4-hydroxybutyl]-N-methylamine (24.0 g) as a pale yellow oil; NMR (CDCl₃): 7.38 (d, 1, J=8.3), 7.27 (d, 1, J=2.1), 7.02 (dd, 1, J=2.1, 8.3), 3.71 (m, 1), 3.53 (m, 1), 2.82-2.72 (m, 3), 2.45 (s, 3), 1.88 (m, 2); MS (CI): m/z=252((M+1), ³⁷Cl ³⁷Cl), 250((M+1), ³⁷Cl ³⁵Cl), 248((M+1), ³⁵Cl ³⁵Cl).

e. (S)-N-[2-(3,4-Dichlorophenyl)-4-hydroxybutyl]-N-methylbenzamide. A solution of (S)-N-[2-(3,4-dichlorophenyl)-4-hydroxybutyl]-N-methylamine (24.0 g) and triethylamine (25.0 g) in dichloromethane (400 mL) was cooled to 0 °C and was treated dropwise with a solution benzoyl chloride (14.05 g) in dichloromethane (100 mL). The mixture was allowed to warm to room temperature, stirred for 16 hours, washed (1N hydrochloric acid (2x), saturated aqueous sodium bicarbonate (2x), and brine), dried, filtered and evaporated to leave a clear oil which solidified on standing. Purification by chromatography (dichloromethane; 99:1 dichloromethane:methanol; 97.5:2.5 dichloromethane:methanol) afforded the benzamide (20.24 g) as a white powder; mp 123-125 °C; [α]_D²⁰ = -18.7 (c=2.03, MeOH); NMR (CD₃OD): 7.53-6.90 (m, 8), 3.84 (m, 1), 3.69 (m, 1), 3.51 (m, 1), 3.42-3.23 (m, 2), 3.05-2.77 (2s, 3), 1.9-1.6 (m, 2); MS (CI): m/z=356((M+1), ³⁷Cl ³⁷Cl), 354((H+1), ³⁷Cl ³⁵Cl), 352((M+1), ³⁵Cl ³⁵Cl).

f. (S)-N-[2-(3,4-Dichlorophenyl)-3-formylpropyl]-N-methylbenzamide. Using a procedure similar to that described in Example 1.1., except using (S)-N-[2-(3,4-dichlorophenyl)-4-hydroxybutyl]-N-methylbenzamide, the aldehyde was isolated as a pale amber syrup; MS (CI): m/z=352((M+1), ³⁷Cl), 350((M+1), ³⁵Cl).

g. (S)-N-[2-(3,4-Dichlorophenyl)-4-[4-(N-methoxy-N-methylcarbamoyl)piperidino]butyl]-N-methylbenzamide. Using a procedure similar to that described in Example 3.b., except using (S)-N-[2-(3,4-dichlorophenyl)-3-propyl]-N-methylbenzamide as the aldehyde component in the reductive alkylation, (S)-N-[2-(3,4-dichlorophenyl)-4-[4-(N-methoxy-N-methylcarbamoyl)piperidino]butyl]-N-methylbenzamide was isolated as a white foam; NMR: 7.56-7.50 (m, 4); 7.41-7.28 (m, 4), 4.22 (s, 1), 3.56 (s, 3), 3.60-3.40 (m, 4), 3.20 (s, 3), 3.30-3.00 (m, 2), 2.50 (m, 1), 2.30 (m, 1), 2.15 (m, 1), 1.97 (m, 2); MS: m/z=451(M).

Example 14. (S)-N-[2-(3,4-Dichlorophenyl)-4-[4-((Z)-1-oximinopropyl)piperidino]butyl]-N-methylbenzamide hydrochloride.

A solution of (S)-N-[2-(3,4-dichlorophenyl)-4-(4-propionylpiperidino)butyl]-N-methylbenzamide (485 mg), hydroxyamine hydrochloride (142 mg) and sodium acetate (167 mg) in acetic acid (8 mL) was stirred for 1 hour at room temperature, was diluted with water (20 mL) and neutralized to pH 7 by addition of sodium bicarbonate. The resulting

mixture was extracted with diethyl ether (3x). The organic extracts were combined, washed successively with saturated aqueous sodium bicarbonate and brine, dried, filtered and evaporated to leave a white foam (496 mg). Purification by chromatography (65:35:10:0.5 dichloromethane:acetone:methanol:ammonium hydroxide) afforded the Z oxime isomer ($R_f=0.61$) as a white foam (97 mg). Treatment of this product with hydrogen chloride (g) in diethyl ether yielded the title compound as a white solid, mp 152-155 °C; NMR: 7.62-7.56 (m,2), 7.43-7.33 (br m,3), 7.23-6.97 (br m,3), 3.85-3.74 (m,2), 3.70-3.45 (m,2), 3.28 (m,1), 3.13-2.80 (m,4), 3.07,2.77 (2s, NCH₃), 2.34 (m,2), 2.23-1.75 (br m,4), 1.09 (m,3); MS (CI): $m/z=492((M+1), ^{37}\text{Cl})$, $490((M+1), ^{35}\text{Cl})$. Analysis for $\text{C}_{26}\text{H}_{32}\text{Cl}_2\text{N}_3\text{O}_2 \cdot 1.0 \text{ HCl} \cdot 0.75 \text{ H}_2\text{O}$: Calculated: C, 57.78; H, 6.62; N, 7.79; Found: C, 57.77; H, 6.40; N, 7.75.

Example 15. (S)-N-[2-(3,4-Dichlorophenyl)-4-[4-((E)-1-oximinopropyl)piperidinolbutyl]-N-methylbenzamide hydrochloride.

From the chromatographic purification in Example 14, a second fraction ($R_f=0.53$) was isolated, which afforded the E oxime isomer as a white foam (172 mg). Treatment of this product with hydrogen chloride (g) in diethyl ether yielded the title compound as a white solid, mp 127-130 °C; NMR: 7.58 (m,2), 7.42-7.31 (br m,3), 7.19-6.95 (br m,3), 3.85-3.74 (m,2), 3.70-3.45 (m,2), 3.24 (m,1), 3.19-2.75 (m,4), 3.04,2.73 (2s, NCH₃), 2.30 (q,2, J=7.6), 2.19-1.68 (br m,4), 1.05 (t,3, J=7.6); MS (CI): $m/z=492((M+1), ^{37}\text{Cl})$, $490((M+1), ^{35}\text{Cl})$. Analysis for $\text{C}_{26}\text{H}_{32}\text{Cl}_2\text{N}_3\text{O}_2 \cdot 1.0 \text{ HCl} \cdot 0.25 \text{ H}_2\text{O}$: Calculated: C, 58.87; H, 6.37; N, 7.92; Found: C, 58.81; H, 6.48; N, 7.83.

Example 16. (S)-N-[2-(3,4-Dichlorophenyl)-4-[4-(1-ethyl-1-hydroxypropyl)piperidino]butyl]-N-methylbenzamide hydrochloride.

Using a procedure similar to that described in Example 1, except using 4-(1-ethyl-1-hydroxypropyl)piperidine, prepared as described in Example 8, and the (S)-aldehyde, prepared as described in Example 13.f., the title compound was isolated as an off white powder, mp 103-105 °C; NMR: 7.61-7.30 (br m,5), 7.20-6.93 (br m,3), 3.86-3.70 (m,2), 3.56 (m,2), 3.25 (m,1), 3.20-2.76 (m,4), 3.05,2.74 (2s, NCH₃), 2.15 (brs,1), 1.90 (brs,3), 1.71-1.38 (m,7), 0.85 (t,6, J=7.3); MS (CI): $m/z=507((M+1), ^{37}\text{Cl})$, $505((M+1), ^{35}\text{Cl})$. Analysis for $\text{C}_{28}\text{H}_{38}\text{Cl}_2\text{N}_2\text{O}_2 \cdot 1.0 \text{ HCl} \cdot 0.50 \text{ H}_2\text{O}$: Calculated: C, 61.04; H, 7.32; N, 5.08; Found: C, 61.18; H, 7.33; N, 5.06.

Example 17. N-[2-(3,4-Dichlorophenyl)-4-(4-propylpiperidino)-butyl]-N-methylbenzamide hydrochloride.

Using a procedure similar to that described in Example 1, except substituting 4-propylpiperidine for the 4-(1-hydroxyethyl)piperidine used therein, the title compound was prepared. Chromatography with dichloromethane:methanol followed by conversion to the hydrochloride salt gave a white solid; mp 73-87 °C; NMR (CD₃OD): 0.9 (t,3, J=7), 1.2-1.6 (m,7), 1.7-2.3 (m,4), 2.5-3.4 (m,8), 3.4-3.6 (m,2), 3.6-3.8 (m,2), 6.8-7.2 (m,3), 7.3-7.4 (m,3), 7.4-7.6 (m,2); MS (CI): $m/z=461(M+1)$. Analysis for: $\text{C}_{26}\text{H}_{34}\text{Cl}_2\text{N}_2\text{O} \cdot 1.0 \text{ HCl} \cdot 0.5 \text{ H}_2\text{O}$: Calculated: C, 61.60; H, 7.16; N, 5.53; Found: C, 61.31; H, 6.96; N, 5.52.

Example 18. N-[2-(3,4-Dichlorophenyl)-4-(4-ethoxycarbonylpiperidino)butyl]-N-methylbenzamide hydrochloride.

Using a procedure similar to that described in Example 1 except substituting 4-ethoxycarbonylpiperidine for the 4-(1-hydroxyethyl)piperidine used therein, the title compound was prepared. Chromatography with dichloromethane:methanol as the eluent, followed by conversion to the hydrochloride salt gave a white solid; mp 48-65; NMR (CD₃OD): 1.1-1.2 (m,3), 1.8-2.2 (m,7), 2.4-3.8 (m,12), 4.0-4.1 (q,J= 7,2), 6.8-7.2 (m,2), 7.2-7.4 (s,4), 7.4-7.8 (m,2) 10.6 (br,1); MS: $m/z=491(M+1)$; Analysis for: $\text{C}_{26}\text{H}_{32}\text{Cl}_2\text{N}_2\text{O}_3 \cdot 1.0 \text{ HCl} \cdot 1.0 \text{ H}_2\text{O}$: Calculated: C, 57.20; H, 6.46; N, 5.13; Found: C, 56.93; H, 6.17; N, 5.06.

Example 19.

The following illustrates representative pharmaceutical dosages forms which may be used for the therapeutic or prophylactic administration of a compound of formula I, or a pharmaceutically acceptable salt thereof, (hereinafter referred to as 'Compound X'):

EP 0 625 509 B1

(i)

Tablet 1	mg/tablet
'Compound X'	100.0
Lactose	77.5
Povidone	15.0
Croscarmellose sodium	12.0
Microcrystalline cellulose	92.5
Magnesium stearate	3.0
	<u>300.0</u>

(ii)

Tablet 2	mg/tablet
'Compound X'	20.0
Microcrystalline cellulose	410.0
Starch	50.0
Sodium starch glycolate	15.0
Magnesium stearate	5.0
	<u>500.0</u>

(iii)

Capsule	mg/capsule
'Compound X'	10.0
Colloidal silicon dioxide	1.5
Lactose	465.5
Pregelatinized starch	120.0
Magnesium stearate	3.0
	<u>600.0</u>

(iv)

5

10

Aerosol	mg/can
'Compound X'	20.0
Oleic acid	10.0
Trichloromonofluoromethane	5,000.0
Dichlorodifluoromethane	10,000.0
Dichlorotetrafluoroethane	5,000.0

15

It will be appreciated that the above pharmaceutical compositions may be varied according to well-known pharmaceutical techniques to accommodate differing amounts and types of active ingredient 'Compound X'. The aerosol (iv) may be used in conjunction with a standard, metered dose aerosol dispenser.

20

25

30

35


40

45


50

55


Formulae




I




II




III



IV



V



IIa

Scheme I.

5

10

15

20

25

30


35

40


45

50


55




VI



VII




VIII



IX

III

IV

Scheme II.

SEQUENCE LISTING

5

(2) INFORMATION FOR SEQ ID NO:1:

10

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 base pairs

(B) TYPE: nucleic acid

15

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GCGCAAGCTT ATGGG

15

25

(2) INFORMATION FOR SEQ ID NO:2:

30

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 18 base pairs

(B) TYPE: nucleic acid

35

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GTCCCCATAA GCTTGCGC

18


45

Claims

50

1. A compound of formula I

55



wherein

Q is phenyl which may bear one or two substituents independently selected from halo, trifluoromethyl, hydroxy, (1-3C)alkoxy, (1-3C)alkyl and methylenedioxy; or Q is thienyl, imidazolyl, benzo[b]thiophenyl or naphthyl any of which may bear a halo substituent; or Q is biphenyl; or Q is carbon-linked indolyl which may bear a benzyl substituent at the 1-position;

Q¹ is hydrogen or (1-3C)alkyl;

Q² is aryl or heteroaryl, which aryl or heteroaryl radical may bear one or more substituents independently selected from halo, hydroxy, (1-4C)alkoxy or (1-4C)alkyl;


R is (1-8C)alkyl or (3-8C)cycloalkyl which may bear one or more substituents selected from the group consisting of halo, (3-6C)cycloalkyl, cyano, nitro, hydroxy, lower alkoxy, lower acyloxy, aroyl, heteroaroyl, oxo, imino (which may bear a (1-6C)alkyl, (3-6C)cycloalkyl, (1-5C)acyl or aroyl substituent) hydroxyimino (which hydroxyimino may bear a lower alkyl or a phenyl substituent on the oxygen), an amino group of formula NR^AR^B, an amino group of formula NR^CR^D, an amidino group of formula C(=NR^G)NR^ER^F, and a carbamoyl group of formula CON(OR^K)R^L, but excluding any compound wherein a hydroxy and an oxo substituent together form a carboxy group, wherein

An amino group of formula NR^AR^B contains zero to about seven carbon atoms and each of R^A and R^B is independently hydrogen, (1-5C)alkyl or (3-6C)cycloalkyl, or R^A and R^B together with the nitrogen to which they are attached form a pyrrolidino, piperidino, morpholino, thiomorpholino (or its S-oxide) or piperazinyl group (which piperazinyl group may bear a methyl or ethyl group at the 4-position); and wherein R^C is hydrogen or lower alkyl and R^D is lower acyl, aroyl or heteroaroyl; or R^D is a group of formula C(=J)NR^ER^F in which J is oxygen, sulfur, NR^G or CHR^H; and wherein the amino group NR^ER^F contains zero to about seven carbon atoms and each of R^E and R^F is independently hydrogen, (1-5C)alkyl or (3-6C)cycloalkyl, or NR^ER^F together with the nitrogen to which they are attached form a pyrrolidino, piperidino, morpholino, thiomorpholino (or its S-oxide) or piperazinyl group (which piperazinyl group may bear a methyl or ethyl group at the 4-position) or R^E is hydrogen or lower alkyl and R^F together with R^G forms an ethylene or trimethylene group; R^G is hydrogen, lower alkyl or together with R^F forms an ethylene or trimethylene group; R^H is cyano, nitro or SO₂R^J and R^J is lower alkyl or phenyl; R^K and R^L are independently (1-3C)alkyl; and in which a cyclic group which is a substituent on R or formed by substitution on R may bear one or more (1-3C)alkyl groups on carbon as further substituents; and in which any aryl or heteroaryl group which is a part of the group R may bear one or more halo, lower alkyl, lower alkoxy, cyano, trifluoromethyl or nitro substituents; or a pharmaceutically acceptable salt thereof.

2. A compound of formula I as claimed in Claim 1 in which Q is 3,4-dichlorophenyl, Q¹ is hydrogen and Q² is phenyl, or a pharmaceutically acceptable salt thereof.
3. A compound of formula I as claimed in Claim 1 in which Q is 3,4-dichlorophenyl; Q¹ is hydrogen; Q² is phenyl; (1-8C)alkyl is ethyl, propyl, butyl, pentyl, isopropyl, 3-pentyl or 4-heptyl; a substituent on the (1-8C)alkyl group of R is hydroxy, acetoxy, oxo, hydroxyimino, methoxyimino, propylamino or acetamido, or two concurrent substituents on R are an oxo substituent and, together on the same carbon, an amino group of formula NR^AR^B, thereby forming a carbamoyl group.
4. A compound of formula I as claimed in Claim 1 or Claim 2 wherein R selected from 1-hydroxyethyl, 1-hydroxypropyl, 1-hydroxybutyl, 1-hydroxy-1-methylethyl, 1-ethyl-1-hydroxy-propyl, 1-hydroxy-1-propylbutyl, 2-hydroxyethyl, acetyl, propionyl, 1-hydroxyiminoethyl, 1-hydroxyiminopropyl, 1-methoxyiminoethyl, 1-methoxyiminopropyl, 2-acetoxyethyl and 2-acetamidoethyl.
5. A compound of formula I as claimed in Claim 1 which is (S)-N-[2-(3,4-dichlorophenyl)-4-[4-(1-ethyl-1-hydroxypropyl)-piperidino]butyl]-N-methylbenzamide, or a pharmaceutically acceptable salt thereof.


6. A salt as claimed in any one of Claims 1-5 which is made with an acid which provides a physiologically acceptable anion.
7. A pharmaceutical composition comprising a pharmaceutically acceptable diluent or carrier and a compound of formula I, or a pharmaceutically acceptable salt thereof, as claimed in any one of Claims 1-6.
8. A method of making a compound of formula I, or a pharmaceutically acceptable salt thereof, as claimed in any one of Claims 1-6 which is characterized by

(a) Alkylating a corresponding piperidine of formula II




II

with an aldehyde of formula III,



III


by reductive alkylation, or with an alkylating agent of formula IV



IV

in which Y is a leaving group;

(b) For a compound of formula I which is a ketone in which R bears an oxo substituent, condensation of a corresponding compound of formula V



V

in which R¹ contains an N,N-disubstituted carbamoyl group with the required organomagnesium or organolithium reagent;

(c) For a compound of formula I wherein R is a primary or secondary hydroxy, reduction of a corresponding compound of formula I wherein R is oxo, using a conventional reducing agent which does not reduce the benzamide moiety;

(d) For a compound of formula I wherein R is hydroxyimino (or an oxygen-substituted hydroxyimino), condensation of a corresponding compound of formula I wherein R is oxo with hydroxylamine (or the corresponding

oxygen-substituted hydroxylamine);


Whereafter, when it is desired to use a protecting group during all or a portion of an above described process; the protecting group then is removed when the final compound is to be formed;

Whereafter, for any of the above procedures, when a pharmaceutically acceptable salt of a compound of formula I is required, it may be obtained by reacting the compound of formula I with an acid affording a physiologically acceptable counterion or by any other conventional procedure;

and in which each of Q, Q¹, Q² and R, except where more particularly described, has the meaning defined in any one of Claims 1-6.

10 Patentansprüche

1. Verbindung der Formel I:



in der bedeuten:

Q Phenyl, das einen oder zwei Substituenten tragen kann, die unabhängig voneinander unter Halogen, Trifluormethyl, Hydroxy, (1-3C)Alkoxy, (1-3C)Alkyl und Methylendioxy ausgewählt sind, oder Thienyl, Imidazolyl, Benzo[b]thiophenyl oder Naphthyl, die jeweils einen Halogen-Substituenten tragen können, oder Biphenyl, oder durch Kohlenstoff gebundenes Indolyl, das einen Benzyl-Substituenten in Position 1 tragen kann, Q¹ Wasserstoff oder (1-3C)Alkyl,

Q² Aryl oder Heteroaryl, wobei der Aryl- oder Heteroaryl-Rest einen oder mehrere Substituenten tragen kann, die unabhängig voneinander unter Halogen, Hydroxy, (1-4C)Alkoxy oder (1-4C)Alkyl ausgewählt sind,


R (1-8C)Alkyl oder (3-8C)Cycloalkyl, das einen oder mehrere Substituenten tragen kann, die unter Halogen, (3-6C)Cycloalkyl, Cyano, Nitro, Hydroxy, Niederalkoxy, Niederacyloxy, Aroyl, Heteroaroyl, Oxo, Imino (das einen (1-6C)Alkyl-, (3-6C)Cycloalkyl-, (1-5C)Acyl- oder Aroyl-Substituenten tragen kann), Hydroxylimino (welches Hydroxylimino einen Niederalkyl- oder einen Phenyl-Substituenten am Sauerstoff tragen kann), einer Amino-Gruppe der Formel NR^AR^B, einer Amino-Gruppe der Formel NR^CR^D, einer Amidino-Gruppe der Formel C(=NR^G)NR^ER^F und einer Carbamoyl-Gruppe der Formel CON(OR^K)R^L ausgewählt sind, wobei aber jede Verbindung ausgenommen ist, bei der ein Hydroxy- und ein Oxo-Substituent zusammen eine Carboxy-Gruppe bilden, wobei

eine Amino-Gruppe der Formel NR^AR^B 0 bis etwa 7 Kohlenstoffatome enthält und jeder der Substituenten R^A und R^B unabhängig voneinander Wasserstoff, (1-5C)Alkyl oder (3-6C)Cycloalkyl bedeutet, oder R^A und R^B zusammen mit dem Stickstoff, an den sie gebunden sind, eine Pyrrolidino-, Piperidino-, Morpholino-, Thiomorpholino- (oder deren S-Oxid) oder Piperaziny-Gruppe (welche Piperaziny-Gruppe eine Methyl- oder Ethyl-Gruppe in der Position 4 tragen kann) bilden, und wobei R^C Wasserstoff oder Niederalkyl und R^D Niederacyl, Aroyl oder Heteroaroyl ist; oder R^D eine Gruppe der Formel C(=J)NR^ER^F ist, in der J Sauerstoff, Schwefel, NR^G oder CHR^H ist, und wobei die Amino-Gruppe NR^ER^F 0 bis etwa 7 Kohlenstoffatome enthält und die Substituenten R^E und R^F jeweils unabhängig voneinander Wasserstoff, (1-5C)Alkyl oder (3-6C)Cycloalkyl bedeuten, oder NR^ER^F zusammen mit dem Stickstoff, an den sie gebunden sind, eine Pyrrolidino-, Piperidino-, Morpholino-, Thiomorpholino- (oder deren S-Oxid)- oder Piperaziny-Gruppe (welche Piperaziny-Gruppe eine Methyl- oder Ethyl-Gruppe in der Position 4 tragen kann) bilden oder R^E Wasserstoff oder Niederalkyl bedeutet und R^F zusammen mit R^G eine Ethylen- oder Trimethylen-Gruppe bildet; R^G Wasserstoff, Niederalkyl bedeutet oder zusammen mit R^F eine Ethylen- oder Trimethylen-Gruppe bildet, R^H Cyano, Nitro oder SO₂R^J und R^J Niederalkyl oder Phenyl bedeutet, R^K und R^L unabhängig voneinander (1-3C)Alkyl bedeuten, und wobei eine cyclische Gruppe, die ein Substituent an R ist oder durch Substitution an R gebildet wird, eine oder mehrere (1-3C)Alkyl-Gruppen an Kohlenstoff als weitere Substituenten tragen kann und wobei eine beliebige Aryl- oder Heteroaryl-Gruppe, die ein Teil der Gruppe R ist, einen oder mehrere Halogen-, Niederalkyl-, Niederalkoxy-, Cyano-, Trifluormethyl- oder Nitro-Substituenten tragen kann, oder ein pharmazeutisch akzeptables Salz davon.

EP 0 625 509 B1


2. Verbindung der Formel I nach Anspruch 1, wobei Q 3,4-Dichlorphenyl, Q¹ Wasserstoff und Q² Phenyl ist, oder ein pharmazeutisch akzeptables Salz davon.
3. Verbindung der Formel I nach Anspruch 1, in der bedeuten: Q 3,4-Dichlorphenyl, Q¹ Wasserstoff, Q² Phenyl, (1-8C)Alkyl Ethyl, Propyl, Butyl, Pentyl, Isopropyl, 3-Pentyl oder 4-Heptyl, wobei ein Substituent an der (1-8C)Alkyl-Gruppe von R Hydroxy, Acetoxy, Oxo, Hydroxyimino, Methoxyimino, Propylamino oder Acetamido ist, oder wobei zwei sich gleichzeitig an R befindliche Substituenten ein Oxo-Substituent sind und, zusammen am gleichen Kohlenstoff, eine Amino-Gruppe der Formel NR^AR^B, wodurch eine Carbamoyl-Gruppe gebildet wird.
4. Verbindung der Formel I nach Anspruch 1 oder 2, wobei R unter 1-Hydroxyethyl, 1-Hydroxypropyl, 1-Hydroxybutyl, 1-Hydroxy-1-methylethyl, 1-Ethyl-1-hydroxypropyl, 1-Hydroxy-1-propylbutyl, 2-Hydroxyethyl, Acetyl, Propionyl, 1-Hydroxyiminoethyl, 1-Hydroxyiminopropyl, 1-Methoxyiminoethyl, 1-Methoxyiminopropyl, 2-Acetoxyethyl und 2-Acetamidoethyl ausgewählt ist.
5. Verbindung der Formel I nach Anspruch 1, nämlich (S)-N-[2-(3,4-Dichlorphenyl)-4-[4-(1-ethyl-1-hydroxypropyl)piperidino]butyl]-N-methylbenzamid, oder ein pharmazeutisch akzeptables Salz davon.
6. Salz nach einem der Ansprüche 1 bis 5, das mit einer Säure hergestellt wurde, die ein physiologisch akzeptables Anion liefert.
7. Pharmazeutische Zusammensetzung, die ein pharmazeutisch akzeptables Verdünnungsmittel oder Trägermittel und eine Verbindung der Formel I, oder ein pharmazeutisch akzeptables Salz davon, nach einem der Ansprüche 1 bis 6 enthält.
8. Verfahren zur Herstellung einer Verbindung der Formel I, oder eines pharmazeutisch akzeptablen Salzes davon, nach einem der Ansprüche 1 bis 6, das dadurch gekennzeichnet ist, daß:

(a) ein entsprechendes Piperidin der Formel II:




II

mit einem Aldehyd der Formel III:



III


durch reduktive Alkylierung oder mit einem Alkylierungsmittel der Formel IV:



IV

in der Y eine Austrittsgruppe ist, alkyliert wird,

(b) zur Herstellung einer Verbindung der Formel I, die ein Keton ist, bei dem R einen Oxo-Substituenten trägt, eine entsprechende Verbindung der Formel V:



15 in der R¹ eine N,N-disubstituierte Carbamoyl-Gruppe enthält, mit dem erforderlichen Organomagnesium- oder Organolithium-Reagens kondensiert wird,

(c) zur Herstellung einer Verbindung der Formel I, in der R ein primäres oder sekundäres Hydroxy ist, eine entsprechende Verbindung der Formel I, in der R Oxo ist, unter Verwendung eines herkömmlichen Reduktionsmittels, welches den Benzamid-Rest nicht reduziert, reduziert wird,


20 (d) zur Herstellung einer Verbindung der Formel I, in der R Hydroxyimino (oder ein Sauerstoff-substituiertes Hydroxyimino) ist, eine entsprechende Verbindung der Formel I, in der R Oxo ist, mit Hydroxylamin (oder dem entsprechenden Sauerstoff-substituierten Hydroxylamin) kondensiert wird,

worauf, wenn eine Schutzgruppe während des gesamten oder eines Teils des oben beschriebenen Verfahrens verwendet werden soll, die Schutzgruppe entfernt wird, wenn die Endverbindung gebildet werden soll,

25 worauf bei jedem der obigen Verfahren, wenn ein pharmazeutisch akzeptables Salz einer Verbindung der Formel I erforderlich ist, dieses durch Umsetzung der Verbindung der Formel I mit einer Säure, die ein physiologisch akzeptables Gegenion liefert, erhältlich ist, oder mit einem bliebigem anderen herkömmlichen Verfahren, wobei Q, Q¹, Q² und R jeweils die in einem der Ansprüche 1 bis 6 definierten Bedeutungen haben, sofern sie nicht besonders definiert sind.

30 Revendications

1. Composé de formule I



45 dans laquelle

Q représente un radical phényle qui peut porter un ou deux substituants choisis indépendamment entre des substituants halogéno, trifluorométhyle, hydroxy, alkoxy en C₁ à C₃, alkyle en C₁ à C₃ et méthylènedioxy ; ou Q représente un radical thiényle, imidazolyle, benzo[b]thiophényle ou naphtyle, l'un quelconque de ces radicaux pouvant porter un substituant halogéno ; ou Q représente un radical biphénylyle ; ou Q représente un radical indolyte, lié à un atome de carbone, qui peut porter un substituant benzyle en position 1 ;

Q¹ représente l'hydrogène ou un radical alkyle en C₁ à C₃ ;

Q² représente un radical aryle ou hétéroaryle, radical aryle ou hétéroaryle qui peut porter un ou plusieurs substituants choisis indépendamment entre des substituants halogéno, hydroxy, alkoxy en C₁ à C₄ et alkyle en C₁ à C₄ ;

55 R représente un radical alkyle en C₁ à C₈, cycloalkyle en C₃ à C₈ qui peut porter un ou plusieurs substituants choisis dans le groupe consistant en substituants halogéno, cycloalkyle en C₃ à C₆, cyano, nitro, hydroxy, alkoxy inférieur, acyloxy inférieur, aroyle, hétéroaroyle, oxo, imino (pouvant porter un substituant alkyle en C₁ à C₆, cycloalkyle en C₃ à C₆, acyle en C₁ à C₅ ou aroyle), hydroxyimino (radical hydroxyimino pouvant porter un substituant alkyle inférieur ou un substituant phényle sur l'atome d'oxygène), un groupe amino de formule


$\text{NR}^{\text{A}}\text{R}^{\text{B}}$, un groupe amino de formule $\text{NR}^{\text{C}}\text{R}^{\text{D}}$, un groupe amidino de formule $\text{C}(=\text{NR}^{\text{G}})\text{NR}^{\text{E}}\text{R}^{\text{F}}$ et un groupe carbamoyle de formule $\text{CON}(\text{OR}^{\text{K}})\text{R}^{\text{L}}$, mais à l'exclusion de tout composé dans lequel un substituant hydroxy et un substituant ortho forment conjointement un groupe carboxy, dans lequel

un groupe amino de formule $\text{NR}^{\text{A}}\text{R}^{\text{B}}$ contient 0 à environ 7 atomes de carbone et chacun des groupes R^{A} et R^{B} représente indépendamment l'hydrogène, un groupe alkyle en C_1 à C_5 ou cycloalkyle en C_3 à C_6 , ou bien R^{A} et R^{B} , conjointement avec l'atome d'azote auquel ils sont fixés, forment un groupe pyrrolidino, pipéridino, morpholino, thiomorpholino (ou son S-oxyde) ou pipérazinyle (groupe pipérazinyle qui peut porter un groupe méthyle ou éthyle en position 4) ; et dans lequel R^{C} représente l'hydrogène ou un groupe alkyle inférieur et R^{D} représente un groupe acyle inférieur, aroyle ou hétéroaroyle ; ou bien R^{D} représente un groupe de formule $\text{C}(=\text{J})\text{NR}^{\text{E}}\text{R}^{\text{F}}$ dans laquelle J représente l'oxygène, le soufre, un groupe NR^{G} ou CHR^{H} ; et dans lequel le groupe amino $\text{NR}^{\text{E}}\text{R}^{\text{F}}$ contient zéro à environ sept atomes de carbone et chacun des groupes R^{E} et R^{F} représente indépendamment l'hydrogène, un groupe alkyle en C_1 à C_5 ou cycloalkyle en C_3 à C_6 , ou bien $\text{NR}^{\text{E}}\text{R}^{\text{F}}$, conjointement avec l'atome d'azote auquel il est fixé, forme un groupe pyrrolidino, pipéridino, morpholino, thiomorpholino (ou son S-oxyde) ou pipérazinyle (groupe pipérazinyle qui peut porter un groupe méthyle ou éthyle en position 4) ou bien R^{E} représente l'hydrogène ou un groupe alkyle inférieur et R^{F} , conjointement avec R^{G} , forme un groupe éthylène ou triméthylène ; R^{G} représente l'hydrogène, un groupe alkyle inférieur ou, conjointement avec R^{F} , forme un groupe éthylène ou triméthylène ; R^{H} représente un groupe cyano, nitro ou $\text{SO}_2\text{R}^{\text{J}}$ et R^{J} représente un groupe alkyle inférieur ou phényle ; R^{K} et R^{L} représentent indépendamment un groupe alkyle en C_1 à C_3 ; et dans lequel un groupe cyclique qui est un substituant sur R ou qui est formé par substitution sur R peut porter un ou plusieurs groupes alkyle en C_1 à C_3 sur des atomes de carbone, comme substituants supplémentaires ; et dans lequel n'importe quel groupe aroyle ou hétéroaroyle qui fait partie du groupe R peut porter un ou plusieurs substituants halogéno, alkyle inférieur, alkoxy inférieur, cyano, trifluorométhyle ou nitro ;

ou l'un de ses sels pharmaceutiquement acceptables.


2. Composé de formule I suivant la revendication 1, dans lequel Q représente un groupe 3,4-dichlorophényle, Q^1 représente l'hydrogène et Q^2 représente un groupe phényle, ou un de ses sels pharmaceutiquement acceptables.
3. Composé de formule I suivant la revendication 1, dans lequel Q représente un groupe 3,4-dichlorophényle, Q^1 représente l'hydrogène, Q^2 représente un groupe phényle ; le groupe alkyle en C_1 à C_8 est un groupe éthyle, propyle, butyle, pentyle, isopropyle, 3-pentyle ou 4-heptyle ; un substituant sur le groupe alkyle en C_1 à C_8 de R est un substituant hydroxy, acétoxy, oxo, hydroxyimino, méthoxyimino, propylamino ou acétamido, ou bien deux substituants pris conjointement sur R représentent un substituant oxo et, en association sur le même atome de carbone avec un groupe amino de formule $\text{NR}^{\text{A}}\text{R}^{\text{B}}$, forment ainsi un groupe carbamoyle.
4. Composé de formule I suivant la revendication 1 ou la revendication 2, dans lequel R est choisi entre les radicaux 1-hydroxyéthyle, 1-hydroxypropyle, 1-hydroxybutyle, 1-hydroxy-1-méthyléthyle, 1-éthyl-1-hydroxypropyle, 1-hydroxy-1-propylbutyle, 2-hydroxyéthyle, acétyl, propionyle, 1-hydroxyiminoéthyle, 1-hydroxyiminopropyle, 1-méthoxyiminoéthyle, 1-méthoxyiminopropyle, 2-acétoxyéthyle et 2-acétamidoéthyle.
5. Composé de formule I suivant la revendication 1, qui est le (S)-N-[2-(3,4-dichlorophényl)-4-[4-(1-éthyl-1-hydroxypropyl)pipéridino]butyl]-N-méthylbenzamide ou un de ses sels pharmaceutiquement acceptables.
6. Sel suivant l'une quelconque des revendications 1 à 5, qui est préparé avec un acide qui donne un anion physiologiquement acceptable.
7. Composition pharmaceutique comprenant un diluant ou support pharmaceutiquement acceptable et un composé de formule I, ou un de ses sels pharmaceutiquement acceptables, suivant l'une quelconque des revendications 1 à 6.
8. Procédé de préparation d'un composé de formule I, ou d'un de ses sels pharmaceutiquement acceptables, suivant l'une quelconque des revendications 1 à 6, qui est caractérisé par

(a) l'alkylation d'une pipéridine correspondante de formule II




II

avec un aldéhyde de formule III



III


par alkylation réductrice, ou avec un agent d'alkylation de formule IV



IV

dans laquelle Y représente un groupe partant ;

(b) pour un composé de formule I qui est une cétone dans laquelle R porte un substituant oxo, la condensation d'un composé correspondant de formule V



V

dans laquelle R¹ contient un groupe carbamoyle N,N-disubstitué avec le réactif organique de magnésium ou organique de lithium requis ;

(c) pour un composé de formule I dans laquelle R représente un groupe hydroxy primaire ou secondaire, la réduction d'un composé correspondant de formule I dans laquelle R représente un groupe oxo, en utilisant un agent réducteur classique qui ne réduit pas le groupement benzamide ;

(d) pour un composé de formule I dans laquelle R représente un groupe hydroxyimino (ou un groupe hydroxyimino O-substitué), la condensation d'un composé correspondant de formule I dans laquelle R représente un groupe oxo avec l'hydroxylamine (ou l'hydroxylamine O-substituée correspondante) ;

puis, lorsqu'il est désiré d'utiliser un groupe protecteur pendant la totalité ou une partie d'un procédé décrit ci-dessus, l'élimination du groupe protecteur lorsque le composé final doit être formé ;

puis, pour n'importe lequel des modes opératoires précités, lorsqu'un sel pharmaceutiquement acceptable d'un composé de formule I est requis, l'obtention de ce sel par réaction du composé de formule I avec un acide donnant un ion complémentaire physiologiquement acceptable ou par n'importe quel autre mode opératoire classique ;

procédé dans lequel chacun des radicaux Q, Q¹, Q² et R, sauf lorsqu'il fait l'objet d'une description plus particulière, répond à la définition suivant l'une quelconque des revendications 1 à 6.





Fig. 2.

pMEG3/hNK2R CONSTRUCT



*Fig.3.*EXPRESSION OF HUMAN NK-2R IN MEL C88 CELLS